


PRINCE GEORGE'S COUNTY

RESULTS OF THE 1994-1997 MARYLAND BIOLOGICAL STREAM SURVEY: COUNTY ASSESSMENTS



CHESAPEAKE BAY AND
WATERSHED PROGRAMS
MONITORING AND
NON-TIDAL ASSESSMENT
CBWP-MANTA-EA-01-25





Parris N. Glendening
Governor

Kathleen Kennedy Townsend
Lt. Governor

A message to Maryland's citizens

The Maryland Department of Natural Resources (DNR) seeks to preserve, protect and enhance the living resources of the state. Working in partnership with the citizens of Maryland, this worthwhile goal will become a reality. This publication provides information that will increase your understanding of how DNR strives to reach that goal through its many diverse programs.

J. Charles Fox
Secretary

Karen M. White
Deputy Secretary



Maryland Department of Natural Resources
Tawes State Office Building
580 Taylor Avenue
Annapolis, Maryland 21401

Toll free in Maryland: 1-(877) 620 8DNR x8611
Out of state call: 410-260-8611
www.dnr.state.md.us

The facilities and services of the Maryland Department of Natural Resources are available to all without regard to race, color, religion, sex, sexual orientation, age, national origin, physical or mental disability.

This document is available in alternative format upon request from a qualified individual with a disability.

Publication date: December 2001
© MD DNR 2001



PRINTED ON RECYCLED PAPER

PRINCE GEORGE'S COUNTY

Results of the 1994-1997 Maryland Biological Stream Survey: County-Level Assessments

**Christopher J. Millard
Paul F. Kazyak
Anthony P. Prochaska**

December 2001

**Maryland Department of Natural Resources
Resource Assessment Service
Monitoring and Non-Tidal Assessment Division
580 Taylor Avenue
Annapolis, MD 21401**

**THIS PAGE INTENTIONALLY
LEFT BLANK**

FOREWORD

This report is based on results of the Maryland Biological Stream Survey (MBSS), a program funded primarily by the Power Plant Research Program and administered by the Maryland Department of Natural Resources (MDNR). Field data for the MBSS were collected by the Maryland Department of Natural Resources. Analyses of water chemistry samples were conducted by the University of Maryland's Appalachian Laboratory. Much of the initial data analysis was conducted by Versar, Inc. for MDNR's Power Plant Assessment Division.

This report helps fulfill two outcomes in MDNR's Strategic Plan: 1) A Vital and Life Sustaining Chesapeake Bay and Its Tributaries, and 2) Sustainable Populations of Living Resources and Healthy Ecosystems.

ACKNOWLEDGMENTS

The 1994-1997 Maryland Biological Stream Survey has been a cooperative effort among several agencies, consultants and academic institutions. We wish to thank Nancy Roth and Ginny Mercurio from Versar in helping to compile some of the data used in this report. Versar also designed the sampling program, obtained landowners' permissions, and helped manage the data. We are also grateful to the many individuals from Maryland Department of Natural Resources, the University of Maryland's Appalachian Laboratory (AL), and the University of Maryland's Wye Research and Education Center (WREC) who comprised the field crews and did a great job collecting the data. MDNR staff also digitized watersheds and calculated land use data, provided quality assurance, and conducted field crew training. Nancy Roth and her colleagues at Versar developed the fish Index of Biotic Integrity, and Dr. Sam Stribling and his staff at Tetra Tech, Inc. developed the benthic Index of Biotic Integrity. Dr. Ray Morgan of AL and Mr. Lenwood Hall of the WREC supervised additional field crews and developed the Physical Habitat Index, and Dr. Keith Eshleman of AL assisted with analyses of data on acidified streams. Drs. Wayne Starnes and Bob Reynolds of the Smithsonian Institution (reptiles and amphibians), Dr. Rich Raesly of Frostburg State University (fish), Rita Villella of the U.S. Geological Survey Leetown Science Center (mussels), and Michael Naylor of MDNR (aquatic vegetation) provided taxonomic verifications of voucher specimens. The success of the project resulted from the strong efforts of all these groups. Special thanks go to Ron Klauda for his editorial support and Brenda Morgan for her assistance in formatting, editing, and organizing the report.

**THIS PAGE INTENTIONALLY
LEFT BLANK**

Table of Contents

FOREWORD	i
ACKNOWLEDGEMENTS	i
INTRODUCTION	1
METHODS	3
MARYLAND BIOLOGICAL STREAM SURVEY DATA	9
COUNTY SUMMARY	10
LIST OF TABLES	
Table 1. Site information and land use data collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.	11
Table 2. Percent occurrence of fish species collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.	15
Table 3. Tolerance Value, Functional Feeding Group, Habit, and Percent Occurrence of benthic macroinvertebrate taxa collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.	18
Table 4. Percent occurrence of reptile and amphibian species collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.	23
Table 5. Physical habitat data for Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.	24
Table 6. Fish Index of Biotic Integrity (F-IBI), Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI), Family-Level Benthic Macroinvertebrate Index of Biotic Integrity (Fam. IBI), and Physical Habitat Index (PHI) scores at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.	27
Table 7. Water chemistry data collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.	30
LIST OF FIGURES	
Figure 1. Land use in Prince George's County.	13
Figure 2. Location of Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.	14
Figure 3. Stream ecological conditions based on the Fish Index of Biotic Integrity (F-IBI) at Maryland Biological Stream Survey sites in Prince George's County.	17
Figure 4. Stream ecological conditions based on the Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI) at Maryland Biological Stream Survey sites in Prince George's County.	22
Figure 5. Stream ecological conditions based on the Physical Habitat Index (PHI) at Maryland Biological Stream Survey sites in Prince George's County.	29
Figure 6. Nitrate-nitrogen concentrations (mg/L) at Maryland Biological Stream Survey sites in Prince George's County.	32
LITERATURE CITED	33
APPENDIX A	
Summary of the types of data collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.	A-1
APPENDIX B	
Physical habitat condition measured by the Maryland Biological Stream Survey, 1994-1997.	B-1

**THIS PAGE INTENTIONALLY
LEFT BLANK**

INTRODUCTION

This report presents county-level data from the 1994–1997 Maryland Biological Stream Survey (MBSS or the Survey). Previous reports have documented interim results from the 1995 (Roth et al. 1997) and 1996 (Roth et al. 1998a) sample years. In addition, a comprehensive final report was produced to assess the “state of the streams” throughout the state (Roth et al. 1999). All previous MBSS reports have presented information by individual drainage basins. Because there is a recognized need for stream health information at the county level, a series of reports were prepared; this report is part of that series. This introductory section recounts the origin of the Survey and describes its components.

Origin of the MBSS

More than 10 years ago, the Maryland Department of Natural Resources (MDNR) recognized that atmospheric deposition was one of the most important environmental problems resulting from the generation of electric power. To determine the extent of acidification of Maryland streams resulting from acidic deposition, MDNR conducted the Maryland Synoptic Stream Chemistry Survey (MSSCS) in 1987. The MSSCS estimated the number and extent of streams at that time affected by or sensitive to acidification statewide and demonstrated the potential for adverse effects on biota from acidification. However, little direct information was available on the biological responses of Maryland streams to water chemistry conditions. Data that were available could not be used (because of methodological differences and spatial coverage limitations) to compare conditions across regions or watersheds (Tornatore et al. 1992). Neither was it possible to assess the interactions between acidic deposition and other anthropogenic and natural influences (CBRM 1989). For these reasons, in 1993, MDNR created the MBSS to provide comprehensive information on the status of biological resources in Maryland streams and how they are affected by acidic deposition and other cumulative effects of anthropogenic stresses.

Description of the MBSS

The MBSS is intended to help environmental decision-

makers protect and restore the natural resources of Maryland. The primary objectives of the MBSS are:

- to assess the current status of biological resources in Maryland’s non-tidal streams;
- to quantify the extent to which acidic deposition has affected or may be affecting biological resources in the state;
- to examine which other water chemistry, physical habitat, and land use factors are important in explaining the current status of biological resources in streams;
- to compile the first statewide inventory of stream biota;
- to establish a benchmark for long-term monitoring of trends in these biological resources; and
- to target future local-scale assessments and mitigation measures needed to restore degraded biological resources.

In creating the Survey, MDNR implemented a probability-based sampling design as a cost-effective way to characterize statewide stream resources. By randomly selecting sites, the Survey can make quantitative inferences about the characteristics of all 9,258 miles of first-to-third-order, non-tidal streams in Maryland (based on stream length on a 1:250,000-scale base map). MDNR recognized that the utility of these estimates depended on accurately measuring appropriate attributes of streams. The Survey focuses on biology for two reasons: (1) organisms themselves have direct societal value and (2) biological communities integrate stresses over time and are a valuable and cost-effective means of assessing ecological integrity (i.e., the capacity of a resource to sustain its inherent potential).

Fish are an important component of stream integrity and one that also contributes to substantial recreational values. For these reasons, fish communities are a primary focus of the Survey. The Survey collects quantitative data for the calculation of population estimates for individual fish species (both game and nongame). These data can also be used to evaluate

fish community composition, individual fish health, and the geographic distribution of commercially important, rare, or non-indigenous fish species. Benthic (bottom-dwelling) macroinvertebrates are another essential component of streams and they constitute the second principal focus of the Survey. The Survey uses rapid bioassessment procedures for collecting benthic macroinvertebrates; these semi-quantitative methods permit comparisons of relative abundance and community composition, and have proven to be an effective way of assessing biological integrity in streams (Hilsenhoff 1987, Lenat 1988, Plafkin et al. 1989, Kerans and Karr 1994, Resh 1995). The Survey also records the presence of reptiles and amphibians (herpetofauna), freshwater mussels, and aquatic plants (both submerged aquatic vegetation (SAV) and emergent macrophytes). The Survey has established rigorous protocols (Kazyak 1996) for each of these sampling components, as well as training and auditing procedures to assure that data quality objectives are met.

Although the MBSS sampling design and protocols provide exceptional information for characterizing the stream resources in Maryland, designation of degraded areas and identification of likely stresses requires additional activities. Assessing the condition of biological resources (whether they are degraded or not degraded) requires the development of ecological indicators that permit the comparison of sampled segment results to minimally impacted reference conditions (i.e., the biological community expected in watersheds with little or no human-induced impacts). The Survey has used its growing database of information collected with consistent methods and broad coverage across the state to develop and test indicators of individual biological components (Stribling et al. 1998, Roth et al. 1998b) and physical habitat quality (Hall et al. 1999). Each of these indicators consists of multiple metrics using the general approach developed for the Index of Biotic Integrity (IBI) (Karr et al. 1986, Karr 1991) and the Chesapeake Bay Benthic Restoration Goals (Ranasinghe et al. 1994). The fish and benthic macroinvertebrate IBIs (which combine attributes of both the number and the type of species found) are widely accepted indicators that have been adapted for use in a variety of geographic locations (Miller et al. 1988, Cairns and Pratt 1993, Simon 1999). The Survey is investigating the possibility

of developing additional indicators (e.g., amphibians in small streams with few or no fish) and combining components into a composite indicator of biological integrity.

In addition to developing reference-based indicators, the Survey is applying a variety of analytical methods to the question of which stressors are most closely associated with degraded streams. This involves correlational and multivariate analyses of water chemistry, physical habitat, land use, and biological information (e.g., presence of non-native species). The biological information also provides a valuable opportunity for documenting aquatic biodiversity across the state; the distribution and abundance of species previously designated as rare only by anecdotal evidence can be determined, and unique combinations of species at the ecosystem and landscape levels can be identified. Land use and other landscape-scale metrics will play an important role in identifying the relative contributions of different stressors to the cumulative impact on stream resources. Ultimately, the Survey seeks to provide an integrated assessment of the problems facing Maryland streams that will facilitate interdisciplinary solutions for their restoration. The survey also provides resource managers with the locations of relatively undisturbed streams and watersheds that deserve protection.

METHODS

This section presents the specific study design and procedures used to implement the Maryland Biological Stream Survey. The study area of concern and the sampling design developed to characterize it are presented, along with field and laboratory methods for each component: fish, benthic macroinvertebrates, reptiles and amphibians, physical habitat, and water chemistry. Methods for aquatic vegetation and mussel sampling are presented, but the resulting data are not included in this report. A full description of MBSS methods can be found in Kazyak (1996).

MBSS Study Design

The Survey study area comprises 17 distinct drainage basins across the state. Random sampling was used to allow the estimation of unbiased summary statistics (e.g., means, proportions, and their respective variances) for the entire state, a particular basin, and subpopulations of interest (e.g., streams with pH < 5).

Because it would have been cost prohibitive to visit a sufficient number of sites in all basins in a single year, lattice sampling was used to schedule sampling of all basins over a three-year period, 1995-1997. Lattice sampling, also known as multistratification, is a cost-effective means of allocating effort across time in a large geographic area (Heimbuch 1999, Jessen 1978, Cochran 1977). A table, or lattice, was formed by arranging 17 basins in 17 rows, and the years in 3 columns. Lattice sampling was the method used for selecting cells from this 17x3 table so that all basins would be sampled over a three-year period and all basins would have a non-zero probability of being sampled in a given year. The data presented in this report include those collected at random sampling sites within the 17 principal basins in Maryland, as well as sites from the 1994 demonstration project. Because no estimates were calculated for this report, these data were included to supplement the number of sites.

The sampling frame for the Survey was constructed by overlaying basin boundaries on a map of all blue-line stream reaches in the study area as digitized on a U.S. Geological Survey 1:250,000 scale topographic map. This sample frame was similar to that used by the earlier Maryland Synoptic Stream Chemistry Survey

(MSSCS) conducted in 1987 (Knapp and Saunders 1987, Knapp et al. 1988). The Strahler convention (Strahler 1957) was used for ranking stream reaches by order; first-order reaches, for example, are the most upstream reaches in the branching stream system. Sampling was restricted to non-tidal, third-order and smaller stream reaches, excluding impoundments that were non-wadable or that substantially altered the riverine nature of the reach (Kazyak 1994). Together, these first-through third-order streams comprise about 90% of all stream and river miles in Maryland. Stream reaches were further divided into non-overlapping, 75-meter segments; these segments were the elementary sampling units from which biological, water chemistry, and physical habitat data were collected.

The 1995-1997 MBSS study design was based on stratified random sampling of segments within each basin; each basin was stratified by stream order. Within a stream order, the number of segments sampled per basin is proportional to the number of stream miles in the basin. To achieve the target number of samples per stream order within each basin, a given number of segments were randomly selected from each basin and ranked in order of selection. In all basins, extra segments were selected as a contingency against loss of sampling sites from restricted access to selected streams or from streams that were dry, too deep, or otherwise unsampleable owing to field conditions. In some basins, where only a small number of sites would have been selected using this method, additional random sites were selected to increase sample size. These extra sites (selected at random using the method described above) were used to provide better basinwide estimates; they were not included in the estimates of statewide conditions.

Permissions were obtained to access privately owned land adjacent to or near each stream segment. The procedures for obtaining permissions are described in Chaillou (1995). Because landowner permissions were obtained in a synoptic fashion and some variation in these rates occurred, we obtained more permissions than were needed for the Survey. Only the highest ranking sites were sampled until the target goal for that basin was reached. For the three year study, the success rate for obtaining permission to access stream sampling segments was high. Eighty-eight percent of sites that were targeted for permission were sampled.

Reasons for permission denial varied and generally reflected the preferences of landowners regarding property access, rather than any specific types of land. In rare cases, permission denial may affect the interpretation of Survey estimates, but only where denials occur in streams with characteristics that differ from the general population of streams. In one example of potential bias, several sites with known coal mining activities in the North Branch Potomac basin denied permission to sample, likely under representing the proportion of acid mine drainage streams in the population.

Field and Laboratory Methods

Benthic macroinvertebrate and water quality sampling were conducted in spring, when the benthos are thought to be reliable indicators of environmental stress (Plafkin et al. 1989) and when acid deposition effects are often the most pronounced. Fish, reptiles and amphibians, aquatic vegetation, and mussel sampling, along with physical habitat evaluations, were conducted during the low-flow period in summer. Fish community composition tends to be stable during summer, and low flow is advantageous for electrofishing. Because low-flow conditions in summer may be a primary factor limiting the abundance and distribution of fish populations, habitat assessments were performed during the summer. The sample size in summer is lower than in spring because some streams were dry in summer or were, in rare cases, otherwise unsampleable.

To reduce temporal variability, sampling during spring and summer was conducted within specific, relatively narrow time intervals, referred to as index periods (Janicki et al. 1993). These index periods were defined by degree-day limits for specific parts of the state. This approach provided a synoptic assessment of the current status of stream biota, water quality, and physical habitat in the 17 basins sampled. The spring index period was the time period between approximately March 1 and May 1, with end of the index period determined by degree-day accumulation as specified in Hilsenhoff (1987). In reality, most spring samples (78%) were collected in March, well before degree-day accumulation limits were approached. The summer index period was between June 1 and September 30 (Kazyak 1994).

Data Collection and Measurement

Field sampling followed procedures specified in the MBSS sampling manual (e.g., Kazyak 1996). A summary of the variables measured and the field and laboratory methods used to conduct the sampling follows.

Fish

Fish were sampled during the summer index period using double-pass electrofishing within 75-meter stream segments. Block nets were placed at each end of the segment and direct current backpack electrofishing units were used to sample the entire segment. An attempt was made to thoroughly fish each segment, and consistent effort was applied over the two passes. This sampling approach allowed calculation of several metrics useful in calculating a biological index and produced unbiased estimates of fish species abundance.

In small streams, a single electrofishing unit was used. In larger streams, two to five units were employed to effectively sample the site. Captured fish were identified to species, counted, weighed, and released. Any individuals that could not be identified to species were retained for laboratory confirmation. For each pass, all individuals of each gamefish species (defined as trout, bass, walleye, pike, chain pickerel, and striped bass) were measured for total length and examined for visible external pathologies or anomalies. For nongame species, up to 100 fish of each species (from both passes) were examined for visible external pathologies or anomalies. For each pass, all non-game species were weighed together for an aggregate biomass measurement; gamefish were also weighed in aggregate to the nearest 10 g.

Electrofishing was also conducted at supplemental, non-randomly selected sites during the summer index period. The presence of each species of fish was recorded for these segments to provide additional qualitative information on statewide fish distributions. Sampling effort at most qualitative sites was based on doubling the elapsed time since the last species was recorded or a minimum of 600 seconds of electrofishing effort.

After processing the fish collected in the field, voucher

specimens were retained for each species not previously collected in the drainage basin. In addition, all individuals which could not be positively identified in the field were retained. The remaining fish were released. All voucher specimens and fish retained for positive identification in the laboratory were examined and verified by the MBSS Quality Assurance Officer or ichthyologists at Frostburg State University, Frostburg, Maryland or the Smithsonian Institution, Washington, DC.

Benthic Macroinvertebrates

Benthic macroinvertebrates were collected to provide a qualitative description of the community composition at each sampling site (Kazyak 1996). Sampling was conducted during the spring index period. Benthic community data were collected for the purpose of calculating biological metrics, such as those described in EPA's Rapid Bioassessment Protocols (Plafkin et al. 1989), and use as an indicator of biological integrity for Maryland streams.

At each segment, a 600 micron mesh "D" net was used to collect organisms from habitats likely to support the greatest taxonomic diversity. A riffle area was preferred, but other habitats were also sampled using a variety of techniques including kicking, jabbing, and gently rubbing hard surfaces by hand to dislodge organisms. If available, other habitat types were sampled, including rootwads, woody debris, leaf packs, macrophytes, and undercut banks. Each jab covered one square foot, and a total of approximately 2.0 m² (20 square feet) of combined substrates was sampled and preserved in 70% ethanol. In the laboratory, the preserved sample was transferred to a gridded pan and organisms were picked from randomly selected grid cells until the cell that contained the 100th individual (if possible) was completely picked. Some samples had fewer than 100 individuals. The benthic macroinvertebrates were identified to genus, or lowest practicable taxon, in the laboratory.

Index of Biotic Integrity

Sites were evaluated using both the fish (F-IBI) and benthic macroinvertebrate (B-IBI) IBIs developed for the MBSS (for detailed methods, see Roth et al. 1997 and Stribling et al. 1998). IBI scores for the MBSS are

determined by comparing the fish or benthic macroinvertebrate assemblages at each site to those found at minimally impacted reference sites. Three separate formulations were employed for the fish IBI, one for each of three distinct geographic areas: Coastal Plain, Eastern Piedmont, and Highland. The two formulations used for the benthic IBI cover the Coastal Plain and non-Coastal Plain regions. Individual metrics for the IBI are scored 1, 3, or 5, based on comparison with the distribution of metric values at reference sites. For either the individual metrics or total IBI, a score of 3 or greater is considered comparable to reference site conditions, while scores falling below this threshold differ significantly from the reference conditions. Scores for the MBSS IBIs are calculated as the mean of the individual metric scores and therefore range from 1 to 5. Some other programs have used a similar approach (e.g., Weisberg et al. 1997), while others have instead computed the IBI as the total of individual metric scores. For example, Karr et al. (1986) calculated IBI as the sum of 12 metric scores, with totals ranging from 12 to 60 points.

Reptiles and Amphibians

At each sample segment, reptiles and amphibians were identified and the presence of observed species was recorded during the summer index period. A search of the riparian area was conducted within 5 meters of the stream on both sides of the 75-meter segment. Any reptiles and amphibians collected during the electrofishing of the stream segment were also included in the species list. Individuals were identified to species when possible. Voucher specimens and individuals not positively identifiable in the field were retained for examination in the laboratory and confirmation by herpetologists at the Smithsonian Institution, Washington, DC, or Towson University, Towson, Maryland.

Physical Habitat

Habitat assessments were conducted at all stream segments as a means of assessing the importance of physical habitat to the biological integrity and fishability of freshwater streams in Maryland. Procedures for habitat assessments (Kazyak 1996) were derived from two currently used methodologies: EPA's Rapid

Bioassessment Protocols (RBPs) (Plafkin et al. 1989), as modified by Barbour and Stribling (1991), and the Ohio EPA's Qualitative Habitat Evaluation Index (QHEI) (Ohio EPA 1987, Rankin 1989). A number of characteristics (instream habitat, epifaunal substrate, velocity/depth diversity, pool/glide/eddy quality, riffle/run quality, channel alteration, bank stability, embeddedness, channel flow status, and shading) were assessed qualitatively, based on visual observations within each 75-meter sample segment. Riparian zone vegetation width was estimated to the nearest meter, up to 50 meters from the stream. Additional observations of the surrounding area were used to assign ratings for aesthetic value (based on visible signs of human refuse at a site) and remoteness (based on distance from the nearest road, accessibility, and evidence of human activity). Also recorded were the presence or absence of various stream features including substrate types, various morphological characteristics, beaver ponds, point sources, and stream channelization. Local land uses visible from the stream segment and riparian vegetation type were also noted. Several additional physical characteristics were measured quantitatively to further characterize the habitat for each segment (see Kazyak 1996 for details). Quantitative measurements of the segment included maximum depth, stream gradient, velocity, thalweg depth, number of functional rootwads, number of functional large woody debris, wetted width, sinuosity, and overbank flood height. A velocity/depth profile was measured or other data were collected to enable calculation of discharge.

Physical Habitat Index

The Physical Habitat Index (PHI) was developed using MBSS data from 1994 to 1997 (Hall et al. 1999). As was the case in development of the fish and benthic IBIs, the conceptual approach was based on evaluating the relative importance (discriminatory power) of individual metrics and combinations of metrics explaining natural differences in streams throughout Maryland. These metrics were derived from both quantitative and qualitative habitat data collected during the summer index period. Based on analyses conducted for both fish IBI (Roth et al. 1998) and benthic macroinvertebrate IBI (Stribling et al. 1998) development in Maryland, the State was divided into two regions: the Coastal Plain and non-Coastal Plain.

The resulting index was then adjusted to a centile scale that rated each sample segment as follows: Good - 72 to 100; Fair - 42 to 71.9; Poor - 12 to 41.9; and Very Poor - 0 to 11.9.

Water Chemistry

During the spring index period, water samples were collected at each site for analysis of pH, acid neutralizing capacity (ANC), conductivity, sulfate, nitrate-nitrogen, and dissolved organic carbon (DOC). These variables describe basic water quality conditions with an emphasis on factors related to acidic deposition.

Grab samples were collected in one-liter bottles for analysis of all analytes except pH. Water samples for pH were collected with 60 ml syringes, which allowed purging of air bubbles to minimize changes in carbon dioxide content (EPA 1987). Samples were stored on wet ice and shipped on wet ice to the analytical laboratory within 48 hours. Laboratory analyses were carried out by the University of Maryland's Appalachian Laboratory in Frostburg.

Chemical analysis of water samples followed standard methods described in EPA's Handbook of Methods for Acid Deposition Studies (EPA 1987). EPA protocols were followed, except that ANC sample volume was reduced to 40 ml to ease handling. Routine daily quality control (QC) checks included processing duplicate, blank, and calibration samples according to EPA guidelines for each analyte. Field duplicates were taken at 5% of all sites. Routine QC checks helped to identify and correct errors in sampling routines or instrumentation at the earliest possible stage.

During the summer index period, in situ measurements of dissolved oxygen (DO), pH, temperature, and conductivity were collected at each site to further characterize existing water quality conditions that might influence biological communities. Measurements were made at an undisturbed section of the segment, usually in the middle of the stream channel, using electrode probes. Instruments were calibrated daily and calibration logbooks were maintained to document instrument performance.

Recognizing that water temperature is an important factor affecting stream condition, but one that varies

daily and seasonally, temperature loggers were deployed at 220 sites in five basins during 1997. The basins sampled were: the Choptank, Susquehanna, Potomac Washington Metro, Patuxent, and Pocomoke. Onset Computer Corporation Optic Stowaway temperature loggers were anchored in each site during the summer index period. Water temperature was recorded every 15 minutes from June 15 until mid-September.

Mussels

During the summer index period, freshwater mussels were sampled qualitatively by examining each 75-meter stream segment for their presence. Mussels were identified to species, their presence recorded, and subsequently released. Species not positively identifiable in the field were retained for confirmation by U.S. Geological Survey (USGS) Biological Resources Division staff.

Aquatic Vegetation

Aquatic vegetation was sampled qualitatively by examining each 75-meter segment for the presence of aquatic plants. Plants were identified to species and their presence recorded for each site. While the primary objective was to document the presence of submerged aquatic vegetation (SAV), emergent and floating aquatic vegetation was also recorded when encountered. Species not positively identifiable in the field were retained for laboratory examination and confirmation by MDNR's staff expert on SAV. Due to the difficulty in long-term preservation, no permanent vouchers of aquatic vegetation were retained.

Data Management

All crews used standardized pre-printed data forms developed for the Survey to ensure that all data for each sampling segment were recorded and standard units of measure were used (Kazyak 1996). Using standard data forms facilitated data entry and minimized transcription error. The field crew leader and a second reviewer checked all data sheets for completeness and legibility before leaving each sampling location. Original data sheets were sent to the Data Management Officer for further review and data entry, while copies were retained by the field crews.

A custom database application, in which the input module was designed to match each of the field data sheets, was used for data entry. Data were independently entered into two databases and compared using a computer program as a quality-control procedure. Differences between the two databases were resolved from original data sheets or through discussions with field crew leaders.

**THIS PAGE INTENTIONALLY
LEFT BLANK**

Maryland Biological Stream Survey Data

COUNTY SUMMARY

A total of 83 quantitative sites were sampled in Prince George's county by MBSS sampling crews from 1994–1997 (Table 1; Figure 2). Qualitative fish sampling was conducted at an additional 14 sites to provide a more complete picture of fish species distributions. Appendix A provides a summary of the types of data available for each of the sites sampled.

Species Highlights

Fifty-seven fish species were collected in the small to mid-sized streams that were sampled; this number ranks first among the counties in the state. Of the sites sampled, 5% contained no fish (Table 2). Possible reasons for the elevated number of fishless sites include migration barriers from road culverts, reduced baseflow and greatly increased stormflow from impervious surfaces, and channel modifications that reduce or effectively eliminate fish habitat (e.g., concrete trapezoids). The 187 genera of benthic macroinvertebrates found in Prince George's county ranks this area fifth best among Maryland counties for benthic diversity (Table 3).

A likely contributing factor for both fish and benthic diversity is the proximity of many county streams to the fall line and the mainstem Potomac River with its diverse array of species. This is apparent in the frequency of occurrence of fishes; only American eel was found at more than two thirds of the sites sampled. In many cases, the Potomac may provide a source of re-colonization after episodes of adverse conditions that are common in urban stream systems.

The most commonly encountered fish species in Prince George's county during the 1994–1997 MBSS included American eel, swallowtail shiner, blacknose dace, eastern mudminnow, white sucker, pumpkinseed, and tessellated darter (Table 2). Rare fish species collected include American brook lamprey, comely shiner, banded sunfish, stripeback darter, and glassy darter. Of these, American brook lamprey and stripeback darter are restricted in Maryland to the Patuxent River basin.

Twenty-six different species of reptiles and amphibians were found in or near Prince George's county streams

(Table 4), ranking the county first in the state. No state or federally listed reptiles or amphibians were collected.

Ecological Health

The overall ecological health of Prince George's county's headwater streams can best be described as Fair to Poor. The average F-IBI score among sites was 3.28 (low end of the Fair range, ranking twelfth among counties in the state) and the average B-IBI score among sites was 2.1 (low end of the Poor category, ranking twenty-first among counties in the state). Based on F-IBI and B-IBI scores from individual sites, the highest rated streams in the county are Swanson Creek and Piscatway Creek (Table 5). The worst rated streams include Oxon Run, Indian Creek, Collington Branch, an unnamed tributary to Beaverdam Creek, and an unnamed tributary to Southwest Branch.

Physical Habitat

Physical habitat in Prince George's County was rated as Fair by the Physical Habitat Index. Values ranged from 2.6 to 96.7, with an average score of 57.8 (mid range of the Fair category, ranking ninth among counties in the state) (Table 6; Figure 5). Other noteworthy points about Prince George's county streams include a ranking of sixteenth for both bank stability and instream rootwads (trees whose roots protect banks from erosion and provide habitat for aquatic life). Prince George's county also had the second poorest rating for trash in the state.

Nitrate-Nitrogen

Nitrate-nitrogen values at sites sampled in Prince George's averaged 1.83 mg/L, ranking the county ninth best in the state. The streams with the lowest nitrate values were Mattawoman Creek, Beaverdam Creek, and an unnamed tributary to Summerville Creek (Table 7). The stream with the highest nitrate value was a first-order section of Zekiah Swamp Run. In this stream, the nitrate level was more than five times the EPA limit for drinking water (10 mg/L).

Table 1. Site information and land use data collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997. Basin abbreviations are as follows: LP - Lower Potomac River; PW - Potomac Washington Metro; PX - Patuxent River.

Site	Latitude	Longitude	Stream Name	Basin	Order	Catchment Acres	% Urban	% Agric.	% Forest
CH-S-139-116-95	38.6370	76.7892	Zekiah Swamp Run	LP	1	1664.10	5.51	23.36	63.28
CH-S-139-118-95	38.6660	76.7898	Zekiah Swamp Run	LP	1	70.27	45.15	19.89	29.13
CH-S-231-202-97	38.5740	76.7430	Swanson Cr	PX	2	5479.94	7.73	20.83	63.47
CH-S-231-209-97	38.5970	76.7460	Swanson Cr	PX	2	3791.33	8.65	22.76	59.47
PG-N-003-2-94	38.8983	76.8019	Western Br	PX	2	12343.70	34.45	26.30	34.30
PG-N-003-3-94	38.8987	76.8026	Western Br	PX	2	12337.40	34.44	26.30	34.31
PG-N-007-127-97	39.1080	76.8960	Un Trib To Patuxent R	PX	1	220.78	32.87	9.17	57.34
PG-N-015-1-94	39.0433	76.8971	Indian Cr	PW	3	2570.10	13.90	31.99	36.32
PG-N-015-2-94	39.0444	76.8960	Indian Cr	PW	3	2546.60	13.91	31.65	36.56
PG-N-027-213-97	38.9220	76.8220	Bald Hill Br	PX	2	10239.09	37.56	22.56	35.59
PG-N-028-301-97	39.0210	76.8560	Beaverdam Cr	PW	3	4187.87	15.45	17.47	59.70
PG-N-041-305-97	38.8870	76.8440	Southwest Br	PX	3	6978.36	42.27	25.79	30.33
PG-N-063-2-94	38.8658	76.9320	Oxon Run	PW	1	841.90	70.45	8.09	20.37
PG-N-063-6-94	38.8484	76.9640	Oxon Run	PW	1	3032.60	64.90	12.36	22.43
PG-N-065-103-97	38.9230	76.8860	Un Trib To Beaverdam Cr	PW	1	1867.43	58.81	16.78	23.61
PG-N-068-125-97	38.7330	76.8660	Un Trib To Piscataway Cr	PW	1	1671.47	9.81	28.28	46.50
PG-N-069-1-94	38.6805	76.9864	Piscataway Cr	PW	1	367.60	6.13	42.53	51.35
PG-N-069-2-94	38.6779	76.9856	Piscataway Cr	PW	1	340.30	6.41	45.94	47.65
PG-N-071-212-97	38.8940	76.8480	Un Trib To Southwest Br	PX	2	1364.82	22.81	50.69	23.86
PG-N-081-1-94	39.0472	76.9036	Indian Cr	PW	2	1698.10	16.52	27.83	30.93
PG-N-081-2-94	39.0451	76.9014	Indian Cr	PW	2	1774.60	18.61	27.23	30.53
PG-N-087-115-97	38.9380	76.7570	Un Trib To Collington Br	PX	1	255.50	5.98	57.31	33.91
PG-N-087-2-94	38.9367	76.7541	Collington Br	PX	1	322.50	4.74	51.91	39.78
PG-N-087-3-94	38.9394	76.7598	Collington Br	PX	1	170.80	8.96	59.19	31.85
PG-N-087-4-94	38.9376	76.7572	Collington Br	PX	1	255.80	5.98	57.35	33.89
PG-N-097-121-97	38.9970	76.7690	Horsepen Br	PX	1	3532.69	27.44	23.29	38.53
PG-N-098-320-97	39.0230	76.8730	Beaverdam Cr	PW	3	7728.98	15.77	15.78	61.60
PG-N-117-329-97	39.0010	76.9820	Northwest Br	PW	3	19458.99	29.68	29.57	39.42
PG-N-119-1-94	38.8161	77.0063	Oxon Cr	PW	1	459.80	65.18	7.09	26.47
PG-N-119-2-94	38.8100	76.9915	Oxon Cr	PW	1	158.20	78.70	6.64	11.00
PG-N-125-218-97	38.9210	76.9040	Beaverdam Cr	PW	2	5016.80	68.41	12.05	17.14
PG-N-125-228-97	38.9220	76.9030	Beaverdam Cr	PW	2	5012.32	68.40	12.06	17.15
PG-N-130-327-97	38.7050	76.9750	Piscataway Cr	PW	3	36461.94	23.42	23.00	44.34
PG-N-135-231-97	38.7830	76.7640	Charles Br	PX	2	9013.86	11.02	35.12	51.14
PG-N-137-1-94	39.0406	76.9005	Indian Cr	PW	3	4475.00	17.66	29.42	33.06
PG-N-137-2-94	39.0381	76.9013	Indian Cr	PW	3	4562.40	19.02	28.97	32.49
PG-N-141-1-94	38.8997	76.8041	Western Br	PX	2	12312.60	34.48	26.31	34.31
PG-N-141-215-97	38.8990	76.7970	Northeast Br Western Br	PX	2	5510.39	9.26	53.42	34.14
PG-N-141-223-97	38.8990	76.7950	Northeast Br Western Br	PX	2	5150.74	5.77	56.10	35.00
PG-N-141-2-94	38.8990	76.7996	Northeast Br Western Br	PX	2	5541.10	9.41	53.34	33.98
PG-N-152-124-97	38.7770	76.7770	Southwest Br Charles Br	PX	1	2814.10	15.93	22.41	60.52
PG-N-155-201-97	38.7330	76.8710	Piscataway Cr	PW	2	9903.33	19.15	23.45	42.36
PG-N-163-111-97	39.0250	76.8270	Beaverdam Cr	PW	1	288.14	28.30	37.26	32.43
PG-N-171-309-97	38.9560	76.9750	Northwest Br	PW	3	29984.83	42.03	21.22	35.70
PG-N-172-1-94	38.8108	77.0083	Oxon Cr	PW	2	8549.90	66.23	8.93	24.39
PG-N-172-2-94	38.8140	77.0071	Oxon Cr	PW	2	8377.30	66.60	8.90	24.07
PG-N-190-103-97	38.7750	76.8050	Un Trib To Charles Br	PX	1	656.86	5.37	22.17	72.45
PG-N-194-1-94	38.8843	76.7407	Collington Br	PX	2	7376.00	18.62	44.41	31.98

Table 1 (cont.). Site information and land use data collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997. Basin abbreviations are as follows: LP - Lower Potomac River; PW - Potomac Washington Metro; PX - Patuxent River.

Site	Latitude	Longitude	Stream Name	Basin	Order	Catchment Acres	% Urban	% Agric.	% Forest
PG-N-194-2-94	38.8922	76.7383	Collington Br	PX	2	6699.50	20.09	44.13	30.73
PG-N-201-330-97	38.9640	76.9720	Northwest Br Anacostia R	PW	3	22549.76	34.24	26.66	37.73
PG-N-205-2-94	38.9705	76.8071	Lottsford Br	PX	1	153.10	28.67	28.02	43.31
PG-N-205-4-94	38.9417	76.8130	Lottsford Br	PX	1	1474.30	34.12	31.37	32.36
PG-N-206-1-94	38.9204	76.7831	Northeast Br	PX	1	925.90	5.03	59.16	33.19
PG-N-206-2-94	38.9278	76.7803	Northeast Br	PX	1	704.40	4.78	59.55	32.81
PG-N-213-113-97	38.8890	76.7660	Black Br	PX	1	1053.48	1.72	63.82	31.65
PG-N-216-135-97	38.8960	76.8490	Un Trib To Southwest Br	PX	1	1076.10	25.99	52.65	18.32
PG-N-219-1-94	38.8811	76.8046	Western Br	PX	3	19114.70	26.98	34.07	34.35
PG-N-219-306-97	38.8950	76.8040	Western Br	PX	3	18039.35	26.85	34.47	34.27
PG-N-219-324-97	38.8910	76.8040	Western Br	PX	3	18185.70	26.81	34.27	34.44
PG-N-219-5-94	38.8785	76.8008	Western Br	PX	3	19232.80	26.83	34.15	34.30
PG-N-232-321-97	38.7080	76.9590	Piscataway Cr	PW	3	24814.94	17.28	24.81	48.04
PG-N-239-2-94	39.0495	76.9082	Indian Cr	PW	1	287.10	30.44	30.23	38.63
PG-N-239-3-94	39.0492	76.9074	Indian Cr	PW	1	292.60	29.97	29.67	39.68
PG-N-246-219-97	38.8270	76.9280	Henson Cr	PW	2	5020.91	56.51	12.03	26.34
PG-N-249-128-97	38.7600	76.9320	Pea Hill Br	PW	1	2058.84	54.48	19.19	23.30
PG-N-251-305-97	39.0310	76.9290	Little Paint Br	PW	3	5793.88	40.89	20.70	32.91
PG-N-253-122-97	38.8500	76.8530	Ritchie Br	PX	1	136.36	17.46	17.95	59.85
PG-N-257-303-97	38.7840	76.9850	Henson Cr	PW	3	11641.00	52.30	12.37	31.64
PG-N-257-306-97	38.7940	76.9600	Henson Cr	PW	3	9474.21	55.56	11.55	28.76
PG-N-257-324-97	38.7960	76.9560	Henson Cr	PW	3	9375.61	55.81	11.64	28.41
PG-N-259-1-94	38.8654	76.7487	Collington Br	PX	2	10827.00	13.24	46.90	35.24
PG-N-259-2-94	38.8621	76.7490	Collington Br	PX	2	10859.20	13.20	46.75	35.26
PG-N-260-1-94	39.0517	76.8977	Indian Cr	PW	2	1151.50	5.58	36.02	23.50
PG-N-260-2-94	39.0587	76.8969	Indian Cr	PW	2	937.20	4.81	33.40	21.31
PG-N-271-9-94	38.9555	76.7642	Collington Br	PX	1	192.20	2.16	63.38	29.76
PG-N-274-128-97	38.8900	76.6830	Honey Br	PX	1	1127.35	2.97	68.68	23.17
PG-S-005-220-95	38.6311	77.0387	Mattawoman Creek	LP	2	28038.91	11.40	19.89	53.92
PG-S-007-108-97	38.6100	76.7180	Un Trib To Summerville Cr	PX	1	19.75	0.00	0.00	100.00
PG-S-032-209-95	38.6549	76.8946	Mattawoman Creek	LP	2	8470.54	11.76	20.21	54.38
PG-S-035-301-97	38.5570	76.7390	Swanson Cr	PX	3	12882.01	6.09	22.79	65.47
PG-S-045-317-97	38.5520	76.7200	Swanson Cr	PX	3	14579.23	5.81	22.42	65.49
PG-S-047-211-97	38.6910	76.7400	Rock Br To Spice Cr	PX	2	5323.36	7.35	32.77	57.62
PG-S-052-109-95	38.6528	76.8078	Wolf Den Branch	LP	1	757.17	10.54	15.95	70.32




Figure 1. Land use in Prince George's County (MOP 1994).




Figure 2. Location of Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Table 2. Percent occurrence of fish species collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Family	Common Name	Scientific Name	Number of Occurrences	Percent Occurrence
Petromyzontidae	least brook lamprey	<i>Lampetra aepyptera</i>	16	21.05
	American brook lamprey	<i>Lampetra appendix</i>	10	13.16
	sea lamprey	<i>Petromyzon marinus</i>	6	7.89
Anguillidae	American eel	<i>Anguilla rostrata</i>	56	73.68
Clupeidae	gizzard shad	<i>Dorosoma cepedianum</i>	2	2.63
Cyprinidae	goldfish	<i>Carassius auratus</i>	2	2.63
	rosyside dace	<i>Clinostomus funduloides</i>	34	44.74
	satinfin shiner	<i>Cyprinella analostana</i>	25	32.89
	spotfin shiner	<i>Cyprinella spiloptera</i>	4	5.26
	common carp	<i>Cyprinus carpio</i>	2	2.63
	cutlips minnow	<i>Exoglossum maxillingua</i>	18	23.68
	eastern silvery minnow	<i>Hybognathus regius</i>	1	1.32
	common shiner	<i>Luxilus cornutus</i>	8	10.53
	golden shiner	<i>Notemigonus crysoleucas</i>	23	30.26
	comely shiner	<i>Notropis amoenus</i>	2	2.63
	silverjaw minnow	<i>Notropis buccatus</i>	1	1.32
	spottail shiner	<i>Notropis hudsonius</i>	8	10.53
	swallowtail shiner	<i>Notropis procne</i>	40	52.63
	bluntnose minnow	<i>Pimephales notatus</i>	2	2.63
	blacknose dace	<i>Rhinichthys atratulus</i>	45	59.21
	longnose dace	<i>Rhinichthys cataractae</i>	10	13.16
	creek chub	<i>Semotilus atromaculatus</i>	28	36.84
	fallfish	<i>Semotilus corporalis</i>	28	36.84
Catostomidae	white sucker	<i>Catostomus commersoni</i>	42	55.26
	creek chubsucker	<i>Erimyzon oblongus</i>	32	42.11
	northern hogsucker	<i>Hypentelium nigricans</i>	3	3.95
Ictaluridae	white catfish ^{1,2}	<i>Ameiurus catus</i>		
	yellow bullhead	<i>Ameiurus natalis</i>	15	19.74
	brown bullhead	<i>Ameiurus nebulosus</i>	10	13.16
	channel catfish ^{1,2}	<i>Ictalurus punctatus</i>		
	tadpole madtom	<i>Noturus gyrinus</i>	9	11.84
	marginated madtom	<i>Noturus insignis</i>	7	9.21
Esocidae	redfin pickerel	<i>Esox americanus vermiculatus</i>	17	22.37
	chain pickerel	<i>Esox niger</i>	7	9.21
Umbridae	eastern mudminnow	<i>Umbra pygmaea</i>	44	57.89
Salmonidae	rainbow trout	<i>Oncorhynchus mykiss</i>	1	1.32
Aphredoderidae	pirate perch	<i>Aphredoderus sayanus</i>	7	9.21
Cyprinodontidae	banded killifish	<i>Fundulus diaphanus</i>	5	6.58
	mummichog	<i>Fundulus heteroclitus</i>	4	5.26
Poeciliidae	mosquitofish	<i>Gambusia affinis</i>	3	3.95
Percichthyidae	white perch	<i>Morone americana</i>	1	1.32
	striped bass ¹	<i>Morone saxatilis</i>		
Centrarchidae	bluespotted sunfish	<i>Enneacanthus gloriosus</i>	10	13.16
	banded sunfish	<i>Enneacanthus obesus</i>	1	1.32
	redbreast sunfish	<i>Lepomis auritus</i>	32	42.11
	green sunfish	<i>Lepomis cyanellus</i>	13	17.11
	pumpkinseed	<i>Lepomis gibbosus</i>	41	53.95
	warmouth	<i>Lepomis gulosus</i>	2	2.63
	bluegill	<i>Lepomis macrochirus</i>	30	39.47

Table 2 (cont.). Percent occurrence of fish species collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Family	Common Name	Scientific Name	Number of Occurrences	Percent Occurrence
	smallmouth bass	<i>Micropterus dolomieu</i>	1	1.32
	largemouth bass	<i>Micropterus salmoides</i>	11	14.47
	black crappie	<i>Pomoxis nigromaculatus</i>	2	2.63
Percidae	tessellated darter	<i>Etheostoma olmsted</i>	48	63.16
	glassy darter	<i>Etheostoma vitreum</i>	6	7.89
	yellow perch	<i>Perca flavescens</i>	2	2.63
	stripeback darter	<i>Percina notogramma</i>	7	9.21
	shield darter ²	<i>Percina peltata</i>		
None			4	5.26

¹ Qualitative Sites

² Fourth- or Fifth-Order Sites




Figure 3. Stream ecological conditions based on the Fish Index of Biotic Integrity (F-IBI) at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Table 3. Tolerance Value (TV)¹, Functional Feeding Group (FFG), Habit, and Percent Occurrence of benthic macroinvertebrate taxa² collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997. Abbreviations of habits are as follows: bu - burrower, cn - clinger, cb - climber, sp - sprawler, dv - diver, and sk - skater.

Class	Order	Family	Genus	TV	FFG	Habit	Percent Occurrence
Nematomorpha ³						bu	3.03
Enopla	Hoplonemertea	Tetrastemmatidae	<i>Prostoma</i> Sp.		Predator		16.67
Turbellaria	Tricladida	Planariidae	<i>Cura</i> Sp.			sp	1.52
			<i>Dugesia</i> Sp.	7	Predator	sp	1.52
Oligochaeta	Lumbriculida	Lumbriculidae		10	Collector	bu	28.79
Oligochaeta	Tubificida	Enchytraeidae		10	Collector	bu	15.15
		Naididae		10	Collector	bu	25.76
		Tubificidae		10	Collector	cn	33.33
			<i>Limnodrilus</i> Sp.	10	Collector	cn	18.18
			<i>Spirosperma</i> Sp.	10	Collector	cn	1.52
Hirudinea				8	Predator	sp	3.03
Gastropoda	Basommatophora	Ancylidae	<i>Fissia</i> Sp.	7	Scraper	cb	1.52
		Lymnaeidae	<i>Pseudosuccinea</i> Sp.	6	Collector	cb	1.52
		Physidae	<i>Physella</i> Sp.	8	Scraper	cb	16.67
		Planorbidae	<i>Planorbella</i> Sp.	7	Scraper	cb	1.52
Pelecypoda	Unionoida	Unionidae		6	Filterer	bu	1.52
Pelecypoda	Veneroida	Corbiculidae	<i>Corbicula</i> Sp.	6	Filterer	bu	1.52
		Sphaeriidae			Filterer	bu	1.52
			<i>Pisidium</i> Sp.	8	Filterer	bu	9.09
			<i>Sphaerium</i> Sp.	8	Filterer	bu	7.58
Malacostraca	Amphipoda					sp	4.55
		Crangonyctidae	<i>Crangonyx</i> Sp.	4	Collector	sp	28.79
		Gammaridae	<i>Gammarus</i> Sp.	6	Shredder	sp	15.15
			<i>Stygonectes</i> Sp.	6	Shredder	sp	9.09
		Hyalellidae	<i>Hyalella</i> Sp.	6	Shredder	sp	6.06
Malacostraca	Decapoda	Cambaridae		6	Shredder	sp	1.52
			<i>Orconectes</i> Sp.	6	Shredder	sp	1.52
Malacostraca	Isopoda	Asellidae	<i>Caecidotea</i> Sp.	8	Collector	sp	22.73
Insecta	Collembola						9.09
Insecta	Ephemeroptera	Ameletidae	<i>Ameletus</i> Sp.	0	Collector	sw, cb	1.52
		Baetidae			Collector	sw, cn	4.55
			<i>Acerpenna</i> Sp.	4	Collector	sw, cn	18.18
			<i>Proclon</i> Sp.	4	Collector		1.52
		Caenidae	<i>Caenis</i> Sp.	7	Collector	sp	3.03
		Ephemerellidae	<i>Ephemerella</i> Sp.	2	Collector	cn, sw	12.12
			<i>Eurylophella</i> Sp.	4	Scraper	cn, sp	15.15
		Heptageniidae	<i>Stenonema</i> Sp.	4	Scraper	cn	13.64
		Leptophlebiidae			Collector	sw, cn	1.52
			<i>Leptophlebia</i> Sp.	4	Collector	sw, cn, sp	1.52
			<i>Paraleptophlebia</i> Sp.	2	Collector	sw, cn, sp	3.03
		Siphonuridae	<i>Siphonurus</i> Sp.	7	Collector	sw, cb	1.52
Insecta	Odonata				Predator		1.52
		Aeshnidae	<i>Boyeria</i> Sp.	2	Predator	cb, sp	15.15
		Calopterygidae	<i>Calopteryx</i> Sp.	6	Predator	cb	9.09
		Coenagrionidae			Predator	cb	6.06
			<i>Argia</i> Sp.	8	Predator	cn, cb, sp	10.61
			<i>Ischnura</i> Sp.	9	Predator	cb	1.52
		Cordulegastridae	<i>Cordulegaster</i> Sp.	3	Predator	bu	3.03

Table 3 (cont.). Tolerance Value (TV)¹, Functional Feeding Group (FFG), Habit, and Percent Occurrence of benthic macroinvertebrate taxa² collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997. Abbreviations of habits are as follows: bu - burrower, cn - climber, cb - climber, sp - sprawler, dv - diver, and sk - skater.

Class	Order	Family	Genus	TV	FFG	Habit	Percent Occurrence
Insecta	Plecoptera	Corduliidae	<i>Macromia</i> Sp.	3	Predator	sp	4.55
		Gomphidae			Predator	bu	4.55
			<i>Gomphus</i> Sp.	5	Predator	bu	3.03
		Capniidae	<i>Allocapnia</i> Sp.	3	Shredder	cn	1.52
			<i>Paracapnia</i> Sp.	1	Shredder	-	1.52
		Chloroperlidae			Predator	cn	3.03
			<i>Haploperla</i> Sp.		Predator	cn	3.03
			<i>Sweltsa</i> Sp.		Predator	cn	3.03
		Leuctridae	<i>Leuctra</i> Sp.	0	Shredder	cn	3.03
		Nemouridae			Shredder	sp, cn	10.61
			<i>Amphinemura</i> Sp.	3	Shredder	sp, cn	15.15
			<i>Ostrocerca</i> Sp.		Shredder	sp, cn	3.03
			<i>Prostoia</i> Sp.		Shredder	sp, cn	19.70
		Perlidae	<i>Eccoptura</i> Sp.		Predator	cn	3.03
		Perlodidae			Predator	cn	4.55
			<i>Clioperla</i> Sp.	1	Predator	cn	4.55
			<i>Isoperla</i> Sp.	2	Predator	cn, sp	15.15
		Taeniopterygidae	<i>Oemopteryx</i> Sp.		Shredder	sp, cn	1.52
			<i>Strophopteryx</i> Sp.		Shredder	sp, cn	7.58
Insecta	Hemiptera		<i>Taeniopteryx</i> Sp.	2	Shredder	sp, cn	1.52
		Corixidae			Predator	sw	3.03
Insecta	Megaloptera		<i>Palmarcorixa</i> Sp.		Predator	-	1.52
		Corydalidae	<i>Nigronia</i> Sp.	0	Predator	cn, cb	7.58
Insecta	Trichoptera	Sialidae	<i>Sialis</i> Sp.	4	Predator	bu, cb, cn	3.03
		Glossosomatidae	<i>Glossosoma</i> Sp.	0	Scraper	cn	1.52
		Hydropsychidae	<i>Cheumatopsyche</i> Sp.	5	Filterer	cn	42.42
			<i>Diplectrona</i> Sp.	2	Filterer	cn	9.09
			<i>Hydropsyche</i> Sp.	6	Filterer	cn	43.94
		Lepidostomatidae	<i>Lepidostoma</i> Sp.	3	Shredder	cb, sp, cn	3.03
		Leptoceridae	<i>Oecetis</i> Sp.	8	Predator	cn, sp, cb	1.52
		Limnephilidae			Shredder	cb, sp, cn	13.64
			<i>Hydatophylax</i> Sp.	2	Shredder	sp, cb	3.03
			<i>Ironoquia</i> Sp.	3	Shredder	sp	12.12
			<i>Pycnopsyche</i> Sp.	4	Shredder	sp, cb, cn	7.58
		Philopotamidae	<i>Chimarra</i> Sp.	4	Filterer	cn	3.03
		Phryganeidae	<i>Ptilostomis</i> Sp.	5	Shredder	cb	6.06
		Polycentropodidae	<i>Polycentropus</i> Sp.	5	Filterer	cn	4.55
		Psychomyiidae	<i>Lype</i> Sp.	2	Scraper	cn	6.06
		Rhyacophilidae	<i>Rhyacophila</i> Sp.	1	Predator	cn	3.03
		Uenoidae				cn	1.52
			<i>Neophylax</i> Sp.	3	Scraper	cn	15.15
Insecta	Lepidoptera			6			1.52
Insecta	Coleoptera	Dryopidae	<i>Helichus</i> Sp.	5	Scraper	cn	4.55
		Dytiscidae	<i>Agabus</i> Sp.	5	Predator	sw, dv	1.52
			<i>Hydroporus</i> Sp.	5	Predator	sw, cb	1.52
			<i>Uvarus</i> Sp.	5	Predator	sw, cb	1.52
		Elmidae	<i>Ancyronyx</i> Sp.	2	Scraper	cn, sp	15.15
			<i>Dubiraphia</i> Sp.	6	Scraper	cn, cb	10.61

Table 3 (cont.). Tolerance Value (TV)¹, Functional Feeding Group (FFG), Habit, and Percent Occurrence of benthic macroinvertebrate taxa² collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997. Abbreviations of habits are as follows: bu - burrower, cn - clinger, cb - climber, sp - sprawler, dv - diver, and sk - skater.

Class	Order	Family	Genus	TV	FFG	Habit	Percent Occurrence
Insecta	Diptera	Gyrinidae	<i>Macronychus</i> Sp.	4	Scraper	cn	6.06
			<i>Optioservus</i> Sp.	4	Scraper	cn	13.64
			<i>Oulimnius</i> Sp.	2	Scraper	cn	6.06
			<i>Stenelmis</i> Sp.	6	Scraper	cn	19.70
			<i>Dinentus</i> Sp.	4	Predator	sw, dv	7.58
			<i>Gyrinus</i> Sp.	4	Predator	sw, dv	7.58
		Haliplidae	<i>Peltodytes</i> Sp.	5	Shredder	cb, cn	1.52
		Hydrophilidae	<i>Berosus</i> Sp.	5	Collector	sw, dv, cb	1.52
			<i>Hydrophilus</i> Sp.	5	Collector	sw, dv, cb	1.52
			<i>Tropisternus</i> Sp.	10	Collector	cb	1.52
		Psephenidae	<i>Psephenus</i> Sp.	4	Scraper	cn	1.52
		Ptilodactylidae	<i>Anchytarsus</i> Sp.	4	Shredder	cn	3.03
		Ceratopogonidae					1.52
					Predator	sp, bu	3.03
		Chironomidae	<i>Bezzia</i> Sp.	6	Predator	bu	3.03
			<i>Culicoides</i> Sp.	10	Predator	bu	3.03
			<i>Probezzia</i> Sp.	6	Predator	bu	7.58
			<i>Ablabesmyia</i> Sp.	8	Predator	sp	3.03
			<i>Brillia</i> Sp.	5	Shredder	bu, sp	13.64
			<i>Brundiniella</i> Sp.	5	Predator	bu, sp	1.52
			<i>Cardiocladius</i> Sp.	6	Predator	bu, cn	1.52
			<i>Chironomus</i> Sp.	10	Collector	bu	1.52
			<i>Clinotanytus</i> Sp.	8	Predator	bu	1.52
			<i>Conchapelopia</i> Sp.	6	Predator	sp	28.79
			<i>Corynoneura</i> Sp.	7	Collector	sp	9.09
			<i>Cricotopus</i> Sp.	7	Shredder	cn, bu	25.76
			<i>Cricotopus</i> / <i>Orthocladius</i> Sp.		Shredder		53.03
			<i>Cryptochironomus</i> Sp.	8	Predator	sp, bu	7.58
			<i>Diamesa</i> Sp.	5	Collector	sp	3.03
			<i>Dicrotendipes</i> Sp.	10	Collector	bu	3.03
			<i>Diplocladius</i> Sp.	7	Collector	sp	15.15
			<i>Endochironomus</i> Sp.	10	Shredder	cn	1.52
			<i>Eukiefferiella</i> Sp.	8	Collector	sp	30.30
			<i>Heterotrissocladius</i> Sp.		Collector	sp, bu	3.03
			<i>Hydrobaenus</i> Sp.	8	Scraper	sp	21.21
			<i>Krenopelopia</i> Sp.		Predator	sp	1.52
			<i>Larsia</i> Sp.	6	Predator	sp	7.58
			<i>Meropelopia</i> Sp.	7			1.52
			<i>Micropsectra</i> Sp.	7	Collector	cb, sp	7.58
			<i>Microtendipes</i> Sp.	6	Filterer	cn	3.03
			<i>Nanocladius</i> Sp.	3	Collector	sp	6.06
			<i>Natarsia</i> Sp.	8	Predator	sp	3.03
			<i>Orthocladinae</i> A Sp.		Collector		18.18
			<i>Orthocladius</i> Sp.	6	Collector	sp, bu	10.61
			<i>Paracladopelma</i> Sp.	7	Collector	sp	1.52
			<i>Parametriocnemus</i> Sp.	5	Collector	sp	36.36
			<i>Paratanytarsus</i> Sp.	6	Collector	sp	9.09

Table 3 (cont.). Tolerance Value (TV)¹, Functional Feeding Group (FFG), Habit, and Percent Occurrence of benthic macroinvertebrate taxa² collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997. Abbreviations of habits are as follows: bu - burrower, cn - climber, cb - climber, sp - sprawler, dv - diver, and sk - skater.

Class	Order	Family	Genus	TV	FFG	Habit	Percent Occurrence
			<i>Paratendipes</i> Sp.	8	Collector	bu	1.52
			<i>Pentaneura</i> Sp.	6	Predator	sp	1.52
			<i>Phaenopsectra</i> Sp.	7	Collector	cn	1.52
			<i>Polypedilum</i> Sp.	6	Shredder	cb, cn	33.33
			<i>Procladius</i> Sp.	9	Predator	sp	3.03
			<i>Prodiamesa</i> Sp.	3	Collector	bu, sp	1.52
			<i>Pseudorthocladius</i> Sp.	0	Collector	sp	1.52
			<i>Rheocricotopus</i> Sp.	6	Collector	sp	27.27
			<i>Rheotanytarsus</i> Sp.	6	Filterer	cn	25.76
			<i>Saetheria</i> Sp.	4	Collector	bu	1.52
			<i>Stenochironomus</i> Sp.	5	Shredder	bu	13.64
			<i>Stictochironomus</i> Sp.	9	Collector	bu	4.55
			<i>Sublettea</i> Sp.		Collector	-	1.52
			<i>Symposiocladius</i> Sp.		Predator	sp	9.09
			<i>Sympotthastia</i> Sp.	2	Collector	sp	3.03
			<i>Tanytarsus</i> Sp.	6	Filterer	cb, cn	15.15
			<i>Thienemanniella</i> Sp.	6	Collector	sp	12.12
			<i>Thienemannimyia</i> Sp.		Predator	sp	16.67
			<i>Tribelos</i> Sp.	5	Collector	bu	3.03
			<i>Tvetenia</i> Sp.	5	Collector	sp	3.03
			CHIRONOMINAE	6	Collector		1.52
			ORTHOCLADIINAE		Collector		3.03
			TANYPODINAE		Predator		1.52
			<i>Xylotopus</i> Sp.	2	Shredder	bu	4.55
			<i>Zavrelimyia</i> Sp.	8	Predator	sp	3.03
		Dixidae	<i>Dixa</i> Sp.	4	Predator	sw, cb	1.52
		Empididae	<i>Chelifera</i> Sp.		Predator	sp, bu	13.64
			<i>Clinocera</i> Sp.		Predator	cn	4.55
			<i>Hemerodromia</i> Sp.	6	Predator	sp, bu	33.33
		Psychodidae	<i>Pericoma</i> Sp.	4	Collector		1.52
		Ptychopteridae	<i>Bittacomorpha</i> Sp.	8	Collector	bu	1.52
		Simuliidae		7	Filterer	cn	4.55
			<i>Prosimulium</i> Sp.	7	Filterer	cn	27.27
			<i>Simulium</i> Sp.	7	Filterer	cn	10.61
			<i>Stegopterna</i> Sp.	7	Filterer	cn	30.30
		Tabanidae	<i>Chrysops</i> Sp.	7	Predator	sp, bu	6.06
			<i>Tabanus</i> Sp.	5	Predator	sp, bu	1.52
		Tipulidae	<i>Antocha</i> Sp.	5	Collector	cn	4.55
			<i>Dicranota</i> Sp.	4	Predator	sp, bu	9.09
			<i>Erioptera</i> Sp.	7	Collector	bu	1.52
			<i>Hexatoma</i> Sp.	4	Predator	bu, sp	9.09
			<i>Ormosia</i> Sp.		Collector	bu	4.55
			<i>Pseudolimnophila</i> Sp.	2	Predator	bu	6.06
			<i>Tipula</i> Sp.	4	Shredder	bu	24.24

¹ Tolerance values are on a 0 (extremely sensitive) to 10 (tolerant) scale.

² Taxa not identified to genus are presented in capital letters. Subfamily - Tanypodinae, Orthoclaadiinae, Chironominae.

³ Nematomorpha is a phylum level identification. No further identification was made.




Figure 4. Stream ecological conditions based on the Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI) at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Table 4. Percent occurrence of reptile and amphibian species collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Family	Common Name	Scientific Name	Number of Occurrences	Percent Occurrence
Ambystomatidae	marbled salamander	<i>Ambystoma opacum</i>	1	1.32
Salamandridae	red spotted newt	<i>Notopthalmus v. viridescens</i>	1	1.32
Plethodontidae	eastern mud salamander	<i>Pseudotriton m. montanus</i>	2	2.63
	northern dusky salamander	<i>Desmognathus f. fuscus</i>	9	11.84
	northern two-lined salamander	<i>Eurycea bislineata</i>	21	27.63
Bufonidae	American toad	<i>Bufo americanus</i>	6	7.89
	Fowler's toad	<i>Bufo woodhousii fowleri</i>	4	5.26
Hylidae	northern cricket frog	<i>Acris crepitans</i>	2	2.63
	northern spring peeper	<i>Pseudacris c. crucifer</i>	1	1.32
Ranidae	bullfrog	<i>Rana catesbeiana</i>	25	32.89
	green frog	<i>Rana clamitans melanota</i>	47	61.84
	northern leopard frog	<i>Rana pipiens</i>	2	2.63
	pickerel frog	<i>Rana palaustris</i>	21	27.63
	southern leopard frog	<i>Rana utricularia</i>	3	3.95
	wood frog	<i>Rana sylvatica</i>	2	2.63
Chelydridae	common snapping turtle	<i>Chelydra serpentina</i>	1	1.32
Kinosternidae	common musk turtle	<i>Sternotherus odoratus</i>	5	6.58
Emydidae	eastern box turtle	<i>Terrapene c. carolina</i>	8	10.53
	eastern painted turtle	<i>Chrysemys p. picta</i>	4	5.26
	spotted turtle	<i>Clemmys guttata</i>	1	1.32
Scincidae	five-lined skink	<i>Eumeces fasciatus</i>	1	1.32
Colubridae	black rat snake	<i>Elaphe o. obsoleta</i>	8	10.53
	eastern garter snake	<i>Thamnophis s. sirtalis</i>	1	1.32
	eastern worm snake	<i>Carpophis a. amoenus</i>	1	1.32
	northern ringneck snake	<i>Diadophis punctatus edwardsii</i>	1	1.32
	northern water snake	<i>Nerodia s. sipedon</i>	6	7.89
None			11	14.47

Table 5. Physical habitat data for Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

	Instream Habitat ¹	Velocity/Depth Diversity ¹		Riffle Quality ¹	Percent Shading ¹		Number of Woody Debris		Percent Channel Flow ¹		Bank Stability ¹		Aesthetic Rating ¹		
Site	Epifaunal Substrate ¹		Pool Quality ¹		Percent Embeddedness ¹		Maximum Depth (cm) ¹		Number of Rootwads		Channel Alteration ¹		Riparian Width (m) ¹		
CH-S-139-116-95	6	2	3	7	7	90	50	30	10	0	70	5	17	50	18
CH-S-231-202-97	9	12	11	12	12	20	93	58	4	2	65	7	7	39	17
CH-S-231-209-97	10	9	13	13	11	40	60	63	5	2	70	6	8	50	13
PG-N-003-2-94	17	14	10	15	0	98	65	112	2		95	13	2	0	13
PG-N-003-3-94	14	11	16	16	10	40	95	170	7		90	8	11	50	16
PG-N-007-127-97	1	2	2	2	0	50	60	25	0	0	25	20	19	0	1
PG-N-015-1-94	10	5	11	11	6	40	80	68	4		40	3	3	50	6
PG-N-015-2-94	8	10	11	17	8	40	97	41	4		50	5	4	0	0
PG-N-027-213-97	7	6	11	16	3	100	97	87	13	3	85	5	2	50	7
PG-N-028-301-97	11	4	11	10	7	65	75	83	5	1	50	5	10	50	16
PG-N-041-305-97	10	6	11	18	11	90	90	159	9	2	75	7	7	50	1
PG-N-063-2-94	1	0	1	0	6	99	35	8	0		50	0	19	0	3
PG-N-063-6-94	13	15	10	7	13	65	60	31	0		75	13	11	0	4
PG-N-065-103-97	2	5	5	1	5	0	10	6	0	0	75	0	20	0	2
PG-N-068-125-97	7	1	5	7	4	100	5	38	6	0	20	1	16	50	17
PG-N-069-1-94	15	14	13	9	10	25	85	63	2		35	11	8	0	11
PG-N-069-2-94	16	16	11	11	14	45	55	32	6		40	14	4	50	16
PG-N-071-212-97	11	11	13	14	13	35	80	74	3	1	90	7	7	0	5
PG-N-081-1-94	2	1	1	0	6	100	15	16	0		70	0	19	30	2
PG-N-081-2-94	1	1	1	0	6	100	10	9	0		95	0	19	10	1
PG-N-087-115-97	5	5	7	6	5	95	92	26	3	2	95	17	15	14	17
PG-N-087-2-94	2	2	3	6	0	95	85	80	0		35	0	2	50	16
PG-N-087-4-94	2	2	1	2	0	100	90	11	2		40	3	4	50	19
PG-N-097-121-97	8	4	14	14	11	85	80	90	10	4	60	4	3	50	9
PG-N-098-320-97	16	7	11	16	6	75	80	110	5	0	97	1	18	50	16
PG-N-117-329-97	15	17	11	16	16	30	65	84	1	0	95	5	17	0	10
PG-N-119-1-94	11	12	8	2	6	35	55	39	3		45	10	6	0	2
PG-N-119-2-94	11	11	10	6	5	40	40	51	0		40	3	14	0	2
PG-N-125-218-97	16	3	12	16	5	65	80	64	2	2	100	11	10	0	1
PG-N-125-228-97	16	4	10	18	2	60	85	85	2	0	100	11	12	0	1
PG-N-130-327-97	12	7	13	19	6	35	50	90	3	1	80	5	13	50	9
PG-N-135-231-97	12	5	9	16	11	100	75	130	12	2	80	5	6	50	13
PG-N-137-1-94	6	6	9	7	10	50	10	19	0		95	0	19	0	1
PG-N-137-2-94	11	2	11	17	6	45	15	93	0		20	2	19	3	2
PG-N-141-1-94	8	9	12	13	14	60	90	75	4		85	10	11	0	5
PG-N-141-2-94	5	2	8	5	0	100	80	40	5		95	3	7	0	11

Table 5 (cont.). Physical habitat data for Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

	Instream Habitat ¹	Velocity/Depth Diversity ¹			Riffle Quality ¹	Percent Shading ¹		Number of Woody Debris		Percent Channel Flow ¹		Bank Stability ¹		Aesthetic Rating ¹	
Site	Epifaunal Substrate ¹			Pool Quality ¹		Percent Embeddedness ¹		Maximum Depth (cm) ¹		Number of Rootwads		Channel Alteration ¹		Riparian Width (m) ¹	
PG-N-141-215-97	7	4	8	16	1	90	70	71	6	5	95	8	5	12	7
PG-N-141-223-97	6	5	9	11	11	50	50	45	2	0	99	7	4	0	9
PG-N-152-124-97	7	4	13	13	11	95	70	80	4	4	90	5	4	0	12
PG-N-155-201-97	17	6	15	16	12	35	85	110	7	1	60	8	16	50	16
PG-N-163-111-97	5	3	6	3	6	100	95	16	0	0	95	18	19	50	16
PG-N-171-309-97	15	8	10	18	1	30	20	99	1	0	90	6	17	30	2
PG-N-190-103-97	7	11	11	11	12	25	80	53	4	1	70	9	5	50	15
PG-N-194-1-94	7	11	14	13	13	80	65	99	7		70	5	4	50	18
PG-N-194-2-94	14	6	12	15	9	85	85	150	7		75	4	3	50	10
PG-N-201-330-97	16	16	14	16	15	35	30	107	0	1	100	8	16	12	12
PG-N-205-2-94	3	3	7	4	7	85	90	34	3		90	6	4	50	15
PG-N-205-4-94	4	5	12	11	16	100	75	88	2		95	5	6	0	10
PG-N-206-1-94	9	2	9	1	0	100	50	37	14		100	19	19	50	18
PG-N-206-2-94	1	1	6	0	0	100	85	17	1		55	3	2	0	16
PG-N-213-113-97	2	2	7	6	6	100	95	33	0	0	55	2	1	50	15
PG-N-216-135-97	7	11	9	8	11	35	92	33	1	0	85	7	4	34	5
PG-N-219-1-94	4	2	8	12	0	100	85	121	4		98	2	2	50	10
PG-N-219-306-97	12	10	13	17	7	98	77	90	14	4	89	9	4	50	8
PG-N-219-324-97	14	11	13	18	6	100	80	172	33	2	65	6	4	50	6
PG-N-219-5-94	5	3	12	12	12	100	60	120	13		70	5	3	50	12
PG-N-232-321-97	12	10	15	20	11	45	85	158	7	2	50	14	12	50	17
PG-N-239-2-94	12	11	8	8	8	5	80	28	4		65	8	11	2	6
PG-N-239-3-94	15	11	11	16	10	5	99	53	3		90	15	11	50	10
PG-N-246-219-97	9	10	8	8	6	30	25	43	4	0	75	16	16	50	8
PG-N-251-305-97	11	8	10	7	6	35	70	34	0	0	97	13	18	10	7
PG-N-253-122-97	9	11	10	11	11	25	75	48	0	3	95	8	9	0	7
PG-N-257-303-97	16	11	15	17	12	45	97	102	5	4	70	15	16	50	16
PG-N-257-306-97	16	16	14	17	16	35	85	106	3	2	90	16	16	10	16
PG-N-257-324-97	18	16	11	19	11	40	80	104	2	2	75	11	16	12	16
PG-N-259-1-94	7	2	16	14	11	50	90	130	6		85	5	6	50	10
PG-N-259-2-94	9	5	8	16	3	75	70	120	4		70	5	5	50	12
PG-N-260-1-94	0	0	1	1	0	100	5	2	0		1	0	19	0	0
PG-N-260-2-94	11	2	12	11	12	50	15	108	1		7	5	16	0	12
PG-N-271-9-94	0	0	0	0	0	99	88	4	3		20	1	1	14	15
PG-N-274-128-97	9	11	9	11	9	25	90	48	0	2	40	8	9	50	18
PG-S-007-108-97	3	12	7	9	4	67	82	30	6	1	97	19	16	50	18

Table 5 (cont.). Physical habitat data for Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Site	Instream Habitat ¹	Velocity/Depth Diversity ¹		Pool Quality ¹	Riffle Quality ¹	Percent Shading ¹		Maximum Depth (cm) ¹	Number of Woody Debris	Percent Channel Flow ¹		Bank Stability ¹	Aesthetic Rating ¹		
	Epifaunal Substrate ¹				Percent Embeddedness ¹			Number of Rootwads		Channel Alteration ¹		Riparian Width (m) ¹			
PG-S-032-209-95	6	5	11	14	11	60	75	98	6	0	100	16	14	17	12
PG-S-035-301-97	7	4	13	13	13	100	85	127	4	3	60	4	6	49	11
PG-S-047-211-97	5	4	8	14	0	100	50	107	13	5	99	16	11	50	17
PG-S-052-109-95	11	7	12	11	13	65	90	52	7	0	95	9	6	50	6

¹ MBSS Qualitative Habitat Metric - See Appendix B for Guidance

Table 6. Fish Index of Biotic Integrity (F-IBI), Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI), Family-Level Benthic Macroinvertebrate Index of Biotic Integrity (Fam. IBI), and Physical Habitat Index (PHI) scores at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Site	Stream Name	F-IBI	B-IBI	Fam. IBI	PHI
CH-S-139-116-95	Zekiah Swamp Run	2.25	1.9		13.28
CH-S-139-118-95	Zekiah Swamp Run		1.0		
CH-S-231-202-97	Swanson Cr	4.75	3.3		71.32
CH-S-231-209-97	Swanson Cr	4.75	4.1		72.44
PG-N-003-2-94	Western Br	2.75	1.3	1.29	
PG-N-003-3-94	Western Br	3.25	1.0	1.00	
PG-N-007-127-97	Un Trib To Patuxent R		1.9		2.56
PG-N-015-1-94	Indian Cr	3.50	1.9	1.86	
PG-N-015-2-94	Indian Cr		1.0	1.00	
PG-N-027-213-97	Bald Hill Br	4.50	3.6		50.47
PG-N-028-301-97	Beaverdam Cr	4.00	3.6		65.38
PG-N-041-305-97	Southwest Br	4.00	1.3		75.81
PG-N-063-2-94	Oxon Run	1.00	1.6		
PG-N-063-6-94	Oxon Run	1.50	1.0	1.00	
PG-N-065-103-97	Un Trib To Beaverdam Cr	1.50	1.3		5.16
PG-N-068-125-97	Un Trib To Piscataway Cr	3.50	2.7		16.48
PG-N-069-1-94	Piscataway Cr		3.6	3.57	
PG-N-069-2-94	Piscataway Cr	2.50	4.4		
PG-N-071-212-97	Un Trib To Southwest Br	3.25	2.4		71.55
PG-N-081-1-94	Indian Cr	1.50	1.6		
PG-N-081-2-94	Indian Cr		2.1		
PG-N-087-115-97	Un Trib To Collington Br		1.3		14.06
PG-N-087-2-94	Collington Br	2.00	2.1		
PG-N-087-3-94	Collington Br		1.9	1.86	
PG-N-087-4-94	Collington Br				
PG-N-097-121-97	Horsepen Br	3.50	2.7		63.36
PG-N-098-320-97	Beaverdam Cr	4.00	4.1		88.44
PG-N-117-329-97	Northwest Br	2.33	1.6		85.86
PG-N-119-1-94	Oxon Cr		1.3	1.29	
PG-N-119-2-94	Oxon Cr		1.0	1.00	
PG-N-125-218-97	Beaverdam Cr	3.25	1.6		71.55
PG-N-125-228-97	Beaverdam Cr	2.50	1.3		77.01
PG-N-130-327-97	Piscataway Cr	4.25	3.9		87.87
PG-N-135-231-97	Charles Br	3.75	3.0		76.02
PG-N-137-1-94	Indian Cr	4.00	1.9	1.86	
PG-N-137-2-94	Indian Cr	4.00	1.0	1.00	
PG-N-141-1-94	Western Br	4.25	1.6	1.57	
PG-N-141-2-94	Northeast Br Western Br	4.00	2.7		
PG-N-141-215-97	Northeast Br Western Br	4.75	2.7		40.66
PG-N-141-223-97	Northeast Br Western Br	4.75	2.1		33.97
PG-N-152-124-97	Southwest Br Charles Br	2.75	2.7		54.04
PG-N-155-201-97	Piscataway Cr	4.75	4.1		95.38
PG-N-163-111-97	Beaverdam Cr		1.9		7.79
PG-N-171-309-97	Northwest Br	3.50	1.0		83.73
PG-N-172-1-94	Oxon Cr		2.1	2.14	
PG-N-172-2-94	Oxon Cr		1.0	1.00	
PG-N-190-103-97	Un Trib To Charles Br	1.75	2.4		58.91
PG-N-194-1-94	Collington Br	3.25	2.1		
PG-N-194-2-94	Collington Br	3.75	1.9		

Table 6 (cont.). Fish Index of Biotic Integrity (F-IBI), Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI), Family-Level Benthic Macroinvertebrate Index of Biotic Integrity (Fam. IBI), and Physical Habitat Index (PHI) scores at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Site	Stream Name	F-IBI	B-IBI	Fam. IBI	PHI
PG-N-201-330-97	Northwest Br Anacostia R	3.75	1.0		92.78
PG-N-205-2-94	Lottsford Br	1.00	2.4		
PG-N-205-4-94	Lottsford Br	4.00	1.9		
PG-N-206-1-94	Northeast Br	2.00	1.9		
PG-N-206-2-94	Northeast Br	2.25	2.7		
PG-N-213-113-97	Black Br	2.00	2.1		9.72
PG-N-216-135-97	Un Trib To Southwest Br	1.00	2.1		25.50
PG-N-219-1-94	Western Br		1.0	1.00	
PG-N-219-306-97	Western Br	4.50	3.6		73.31
PG-N-219-324-97	Western Br	4.75	2.7		89.21
PG-N-219-5-94	Western Br	3.75	1.6		
PG-N-232-321-97	Piscataway Cr	3.25	2.4		96.74
PG-N-239-2-94	Indian Cr	3.50	1.9		
PG-N-239-3-94	Indian Cr	2.75	2.4		
PG-N-246-219-97	Henson Cr	3.25	1.9		34.72
PG-N-249-128-97	Pea Hill Br		2.1		
PG-N-251-305-97	Little Paint Br	3.25	1.9		37.51
PG-N-253-122-97	Ritchie Br		2.1		49.36
PG-N-257-303-97	Henson Cr	3.00	1.0		94.42
PG-N-257-306-97	Henson Cr	3.50	1.6		94.65
PG-N-257-324-97	Henson Cr	3.75	2.1		94.81
PG-N-259-1-94	Collington Br	3.75	1.6		
PG-N-259-2-94	Collington Br	3.75	2.1		
PG-N-260-1-94	Indian Cr		1.0	1.00	
PG-N-260-2-94	Indian Cr	3.25	1.0		
PG-N-271-9-94	Collington Br	1.00	1.6		
PG-N-274-128-97	Honey Br	4.00	2.4		61.55
PG-S-005-220-95	Mattawoman Creek		1.6		
PG-S-007-108-97	Un Trib To Summerville Cr		1.3		21.54
PG-S-032-209-95	Mattawoman Creek	2.25	2.4		62.84
PG-S-035-301-97	Swanson Cr	4.75	3.6		63.87
PG-S-045-317-97	Swanson Cr		2.7		
PG-S-047-211-97	Rock Br To Spice Cr	5.00	3.6		50.47
PG-S-052-109-95	Wolf Den Branch	4.00	3.6		49.09




Figure 5. Stream ecological conditions based on the Physical Habitat Index (PHI) at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Table 7. Water chemistry data collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Site	pH	Conductivity (μ S/cm)	Acid Neutralizing Capacity (μ eq/L)	Nitrate (mg/L)	Sulfate (mg/L)	Dissolved Oxygen (mg/L)	Dissolved Organic Carbon (mg/L)
CH-S-139-116-95	5.45	0.073	17.08	0.474	9.185	2.50	5.00
CH-S-139-118-95	6.00	0.702	1715.62	52.866	31.020		120.00
CH-S-231-202-97	6.82	0.086	156.00	0.530	13.730	9.60	2.60
CH-S-231-209-97	6.29	0.076	54.20	0.646	12.302	7.70	3.20
PG-N-003-2-94							
PG-N-003-3-94	6.79	0.196	346.99	0.664	18.898		4.00
PG-N-007-127-97	6.71	0.162	475.10	1.022	18.553	8.70	5.80
PG-N-015-1-94							
PG-N-015-2-94	6.93	0.144	460.30	0.451	18.703		4.00
PG-N-027-213-97	7.03	0.250	566.20	0.742	17.546	9.30	6.60
PG-N-028-301-97	6.69	0.142	209.00	1.525	16.038	7.90	4.00
PG-N-041-305-97	7.25	0.377	978.20	0.712	26.085	6.60	5.70
PG-N-063-2-94							
PG-N-063-6-94	6.99	0.338	650.16	2.084	46.498		3.00
PG-N-065-103-97	7.25	0.354	1226.30	0.813	38.605	5.80	4.00
PG-N-068-125-97	6.75	0.168	278.10	0.596	19.852	6.40	3.00
PG-N-069-1-94	6.51	0.159	111.99	1.153	13.704		2.00
PG-N-069-2-94							
PG-N-071-212-97	7.12	0.473	1005.30	0.548	24.827	7.70	4.70
PG-N-081-1-94							
PG-N-081-2-94	7.05	0.160	551.52	0.632	15.333		4.00
PG-N-087-115-97	6.54	0.271	406.10	0.427	19.009	8.00	6.80
PG-N-087-2-94	6.18	0.213	93.63	0.979	19.300		2.00
PG-N-087-3-94							
PG-N-097-121-97	6.88	0.191	495.80	0.680	17.124	9.00	6.90
PG-N-098-320-97	6.71	0.146	210.50	1.145	15.555	8.50	4.10
PG-N-117-329-97	7.48	0.222	889.50	1.680	13.978	6.90	1.90
PG-N-119-1-94	7.41	0.327	627.17	1.413	50.679		2.00
PG-N-119-2-94							
PG-N-125-218-97	7.30	0.500	1341.10	0.966	30.965	7.50	4.50
PG-N-125-228-97	7.56	0.378	1318.80	0.167	31.498	7.50	3.80
PG-N-130-327-97	6.97	0.175	304.00	0.710	22.855	5.60	2.70
PG-N-135-231-97	6.81	0.180	288.70	0.980	33.647	6.90	2.30
PG-N-137-1-94							
PG-N-137-2-94	6.94	0.174	624.25	0.753	17.743		3.00
PG-N-141-1-94	6.65	0.143	248.67	0.852	21.345		2.00
PG-N-141-2-94							
PG-N-141-215-97	7.06	0.212	382.80	0.792	24.648	6.30	2.50
PG-N-141-223-97	6.99	0.202	339.60	0.824	23.492	8.70	2.30
PG-N-152-124-97	7.00	0.172	409.10	0.780	31.541	7.60	2.60
PG-N-155-201-97	6.92	0.162	320.20	1.008	21.870	6.70	3.80
PG-N-163-111-97	5.91	0.150	182.00	4.708	18.701	7.80	1.80
PG-N-171-309-97	7.63	0.279	1058.90	1.636	17.292	6.50	2.40
PG-N-172-1-94	7.39	0.362	731.32	1.682	55.673		3.00
PG-N-172-2-94							
PG-N-190-103-97	6.37	0.174	110.50	0.712	33.974	9.60	1.80
PG-N-194-1-94	6.75	0.156	292.11	0.734	22.222		3.00
PG-N-194-2-94							

Table 7 (cont.). Water chemistry data collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Site	pH	Conductivity (μ S/cm)	Acid Neutralizing Capacity (μ eq/L)	Nitrate (mg/L)	Sulfate (mg/L)	Dissolved Oxygen (mg/L)	Dissolved Organic Carbon (mg/L)
PG-N-201-330-97	7.72	0.208	883.50	1.410	14.445	8.40	1.90
PG-N-205-2-94							
PG-N-205-4-94	6.70	0.207	337.14	1.186	17.000		2.00
PG-N-206-1-94	6.41	0.133	158.89	0.811	24.194		2.00
PG-N-206-2-94							
PG-N-213-113-97	6.72	0.257	359.00	0.665	19.828	8.60	6.10
PG-N-216-135-97	6.98	0.332	851.90	0.614	26.705	8.30	5.10
PG-N-219-1-94	6.76	0.198	323.96	0.706	20.218		3.00
PG-N-219-306-97	7.14	0.244	531.60	0.566	21.612	7.00	4.00
PG-N-219-324-97	7.06	0.242	527.30	0.672	21.487	5.30	3.80
PG-N-219-5-94	6.76	0.181	320.44	0.693	19.846		3.00
PG-N-232-321-97	7.02	0.156	214.30	0.721	22.191	6.70	2.80
PG-N-239-2-94	7.05	0.181	805.72	1.700	19.910		3.00
PG-N-239-3-94							
PG-N-246-219-97	7.36	0.339	765.30	1.072	24.320	7.00	3.40
PG-N-249-128-97	6.85	0.229	345.50	0.948	19.906		3.10
PG-N-251-305-97	7.64	0.218	592.30	1.192	12.269	9.40	2.30
PG-N-253-122-97	7.23	0.826	2091.40	0.804	326.950	8.40	7.50
PG-N-257-303-97	7.44	0.311	837.80	4.588	14.933	8.60	3.00
PG-N-257-306-97	7.44	0.324	758.20	1.118	26.122	9.40	3.00
PG-N-257-324-97	7.35	0.336	713.70	1.152	26.858	9.40	3.10
PG-N-259-1-94							
PG-N-259-2-94	6.80	0.148	336.29	0.757	21.793		3.00
PG-N-260-1-94	7.23	0.288	980.58	1.670	15.495		7.00
PG-N-260-2-94							
PG-N-271-9-94	6.07	0.235	230.08	4.332	38.647		5.00
PG-N-274-128-97	7.30	0.138	257.20	0.762	20.679	8.70	2.40
PG-S-005-220-95	4.94	0.061	-6.42	0.187	14.716		6.00
PG-S-007-108-97	5.19	0.050	9.50	0.294	8.302	8.70	2.30
PG-S-032-209-95	6.29	0.164	130.69	0.396	13.697	3.20	5.00
PG-S-035-301-97	7.01	0.089	230.40	0.604	14.470	6.90	2.80
PG-S-045-317-97	6.87	0.095	279.30	0.489	15.172		2.70
PG-S-047-211-97	6.61	0.121	137.80	0.753	23.616	6.90	2.20
PG-S-052-109-95	5.98	0.097	58.82	0.705	12.545	6.00	5.00




Figure 6. Nitrate-nitrogen concentrations (mg/L) at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.

Literature Cited

- Barbour, M.T., and J.B. Stribling. 1991. Use of habitat assessment in evaluating the biological integrity of stream communities. In: *Biological Criteria: Research and Regulation*. U.S. Environmental Protection Agency, Washington, D.C. EPA-440/5-91-005. pp. 25-38.
- Cairns, J. and J.R. Pratt. 1993. A history of biological monitoring using benthic macroinvertebrates. In Rosenberg, D.M. and V.H. Resh, eds. *Freshwater Monitoring and Benthic Macroinvertebrates*. Chapman and Hall, New York
- CBRM (Chesapeake Bay Research and Monitoring Division). 1989. Acid deposition in Maryland. Summary of results through 1988. Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division, Annapolis, MD. AD-89-1. NTIS No. PB89-182729/AS.
- Chaillou, J.C. 1995. Technical Memorandum, The Maryland Biological Stream Survey: Summary of landowner contact procedures. Prepared by Versar, Inc., Columbia, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division, Annapolis, MD. CBWP-MANTA-EA-95-1.
- Cochran, W.G. 1977. *Sampling Techniques*. 3rd ed. New York: John Wiley and Sons.
- EPA (U.S. Environmental Protection Agency). 1987. *Handbook of methods for acid deposition studies: Laboratory analysis for surface water chemistry*. EPA-600/4-87/026.
- Hall, L.W., Jr., R.P. Morgan II, E.S. Perry, A. Waltz. 1999. Development of a provisional physical habitat index for Maryland freshwater streams. Prepared by Wye Research and Education Center, Queenstown, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment 455.
- Heimbuch, D.G., J.C. Seibel, H.T. Wilson, P.F. Kazyak. 1999. A multiyear lattice sampling design for Maryland-wide fish abundance estimation. *Journal of Agricultural, Biological, and Environmental Statistics*. 4: 443-455.
- Hilsenhoff, W.L. 1987. An improved biotic index of organic stream pollution. *Great Lakes Entomologist* 20: 31-39.
- Janicki, A., D. Wade, D. Heimbuch, H. Wilson, P. Jacobson, P. Kazyak. 1993. Maryland Biological Stream Survey design report. Prepared for Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division, by Coastal Environmental Services, Linthicum, MD. CBRM-AD-93-1.
- Jessen, R.J. 1978. *Statistical Survey Techniques*. John Wiley, New York.
- Karr, J.R. 1991. Biological integrity: A long-neglected aspect of water resource management. *Ecological Applications* 1: 66-84.
- Karr, J.R., K.D. Fausch, P.L. Angermeier, P.R. Yant, and I.J. Schlosser. 1986. Assessing biological integrity in running waters: a method and its rationale. *Illinois Natural History Survey Special Publication* 5. 28 pp.
- Kazyak, P.F. 1996. Maryland Biological Stream Survey Sampling Manual. Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division.
- Kazyak, P.F. 1994. Maryland Biological Stream Survey Sampling Manual. Prepared by Versar, Inc., Columbia, MD, for Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division.
- Kerans, B.L. and J.R. Karr. 1994. A benthic index of biotic integrity (B-IBI) for rivers of the Tennessee valley. *Ecological Applications* 4: 768-785.
- Knapp, C.M., W.P. Saunders, D.G. Heimbuch, H.S. Greening, and G.J. Filbin. 1988. Maryland

- Synoptic Stream Chemistry Survey: Estimating the number and distribution of streams affected by or at risk from acidification. Prepared by International Science and Technology, Inc., Reston, VA, for the Maryland Department of Natural Resources, Power Plant Research Program, Annapolis, MD. AD-88-2. NTIS No. PB88-213996/AS.
- Knapp, C.M. and W.P. Saunders. 1987. Maryland Synoptic Stream Chemistry Survey design report. Prepared by International Science and Technology, Inc. for Maryland Department of Natural Resources, Power Plant Research Program. AD-87-5.
- Lenat, D.R. 1988. Water quality assessment of streams using a qualitative collection method for benthic macroinvertebrates. *Journal of North American Benthological Society* 7: 222-223.
- Miller, D.L., P.M. Leonard, R.M. Hughes, J.R. Karr, P.B. Moyle, L.H. Schrader, B.A. Thompson, R.A. Daniels, K.S. Fausch, G.A. Fitzhugh, J.R. Gammon, D.B. Halliwell, P.L. Angermeier, and D.J. Orth. 1988. Regional applications of an index of biotic integrity for use in water resource management. *Fisheries* 13(5): 12-20.
- Ohio EPA (Ohio Environmental Protection Agency). 1987. Biological criteria for the protection of aquatic life. Volumes I-III. Ohio EPA, Division of Water Quality Monitoring and Assessment, Surface Water Section, Columbus, Ohio.
- Plafkin, J.L., M.T. Barbour, K.D. Porter, S.K. Gross and R.M. Hughes. 1989. Rapid bioassessment protocols for use in streams and rivers: Benthic macroinvertebrates and fish. U.S. Environmental Protection Agency, Office of Water, Washington, D.C. EPA 440-4-89-001.
- Ranasinghe, J.A., S.B. Weisberg, D.M. Dauer, L.C. Schaffner, R.J. Diaz, and J.B. Frithsen. 1994. Chesapeake Bay Benthic Community Restoration Goals. Prepared by Versar, Inc. for U.S. EPA and the Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division.
- Rankin, E.T. 1989. The Qualitative Habitat Evaluation Index (QHEI): Rationale, methods, and application. Ohio EPA, Division of Water Quality Planning and Assessment, Ecological Analysis Section, Columbus, OH.
- Resh, V. H., 1995. Freshwater benthic macroinvertebrates and rapid assessment procedures for water quality monitoring in the developing and newly industrialized countries. In: Davis, W.S. and T.P. Simon, eds. 1995. Biological assessment and criteria: tools for water resource planning and decision making. Lewis Publishers, Boca Raton, FL.
- Roth, N.E., M.T. Southerland, G. Mercurio, J.C. Chaillou, P.F. Kazyak, S.A. Stranko, A.P. Prochaska, D.G. Heimbuch, and J.C. Seibel. 1999. State of the Streams: 1995-1997 Maryland Biological Stream Survey Results. Prepared by Versar, Inc., Columbia, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division. CBWP-MANTA-EA-99-6.
- Roth, N.E., M.T. Southerland, J.C. Chaillou, G.T. Wilson, D.G. Heimbuch, and J.C. Seibel. 1998a. Maryland Biological Stream Survey: Ecological Status of Non-tidal Streams Sampled in 1996. Prepared by Versar, Inc., Columbia, MD, and Post, Buckley, Schuh, and Jernigan, Inc., Bowie, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division. CBWP-MANTA-EA-97-2.
- Roth, N.E., M.T. Southerland, J.C. Chaillou, R.J. Klauda, P.F. Kazyak, S.A. Stranko, S.B. Weisberg, L.W. Hall, Jr., and R.P. Morgan II. 1998b. Maryland Biological Stream Survey: Development of a Fish Index of Biotic Integrity. *Environmental Management and Assessment* 51: 89-106.
- Roth, N.E., M.T. Southerland, J.C. Chaillou, J.H. Volstad, S.B. Weisberg, H.T. Wilson, D.G. Heimbuch, and J.C. Seibel. 1997. Maryland Biological Stream Survey: Ecological Status of Non-Tidal Streams in Six Basins Sampled in 1995. Prepared by Versar, Inc., Columbia, MD and Coastal Environmental Services, Linthicum, MD for Maryland Department of Natural Resources,

Monitoring and Non-tidal Assessment Division.
CBWP-MANTA-EA-97-2.

Simon, T.P. ed. 1999. Assessing the Sustainability and Biological Integrity of Water Resources Using Fish Communities. CRC Press, Washington DC.

Strahler, A.N. 1957. Quantitative analysis of watershed geomorphology. Transactions of the American Geophysical Union 38(6): 913-920.

Stribling, J.B., B.K. Jessup, J.S. White, D. Boward, M. Hurd. 1998. Development of a benthic index of biotic integrity for Maryland streams. Prepared by Tetra Tech, Inc., Owings Mills, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division, Annapolis, MD. CBWP-EA-98-3.

Tornatore, T., P. Jacobson, J. Gerritsen and A. Janicki. 1992. Data sets for developing measures of biological integrity in Maryland streams. Prepared by Versar, Inc., Columbia, MD, for the Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division, Annapolis, MD. AD-93-7.

Weisberg, S.B., J.A. Ranasinghe, D.M. Dauer, L.C. Schaffner, R.J. Diaz, J.B. Frithsen. 1997. An estuarine benthic index of biotic integrity (B-IBI) for Chesapeake Bay. Estuaries 20: 149-158.

**THIS PAGE INTENTIONALLY
LEFT BLANK**

Appendix A. Summary of the types of data collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997.
Abbreviations used are as follows: F-IBI - Fish Index of Biotic Integrity; B-IBI Benthic Macroinvertebrate Index of Biotic Integrity;
Fam.IBI - Family-Level Benthic Macroinvertebrate Index of Biotic Integrity; PHI - Physical Habitat Index.

<i>Site</i>	<i>Stream Name</i>	<i>Benthic Macroinvertebrate</i>		<i>Habitat</i>		<i>F-IBI</i>		<i>Fam. IBI</i>	
		<i>Fish</i>	<i>Herpetofauna</i>	<i>Water Chemistry</i>			<i>B-IBI</i>		<i>PHI</i>
CH-S-139-116-95	Zekiah Swamp Run	X	X	X	X	X	X		X
CH-S-139-118-95	Zekiah Swamp Run		X			X	X		
CH-S-231-202-97	Swanson Cr	X	X	X	X	X	X		X
CH-S-231-209-97	Swanson Cr	X	X	X	X	X	X		X
PG-N-003-2-94	Western Br	X	X	X		X		X	
PG-N-003-3-94	Western Br	X	X	X	X	X		X	
PG-N-007-127-97	Un Trib To Patuxent R	X	X	X	X		X		X
PG-N-015-1-94	Indian Cr	X	X	X		X		X	
PG-N-015-2-94	Indian Cr	X	X	X	X			X	
PG-N-027-213-97	Bald Hill Br	X	X	X	X	X	X		X
PG-N-028-301-97	Beaverdam Cr	X	X	X	X	X	X		X
PG-N-041-305-97	Southwest Br	X	X	X	X	X	X		X
PG-N-063-2-94	Oxon Run	X	X	X		X	X		
PG-N-063-6-94	Oxon Run	X	X	X	X	X		X	
PG-N-065-103-97	Un Trib To Beaverdam Cr	X	X	X	X	X	X		X
PG-N-068-125-97	Un Trib To Piscataway Cr	X	X	X	X	X	X		X
PG-N-069-1-94	Piscataway Cr	X	X	X	X			X	
PG-N-069-2-94	Piscataway Cr	X	X	X		X	X		
PG-N-071-212-97	Un Trib To Southwest Br	X	X	X	X	X	X		X
PG-N-081-1-94	Indian Cr	X	X	X		X	X		
PG-N-081-2-94	Indian Cr	X	X	X	X		X		
PG-N-087-115-97	Un Trib To Collington Br	X	X	X	X		X		X
PG-N-087-2-94	Collington Br	X	X	X	X	X	X		
PG-N-087-3-94	Collington Br		X					X	
PG-N-087-4-94	Collington Br	X		X	X				
PG-N-097-121-97	Horsepen Br	X	X	X	X	X	X		X
PG-N-098-320-97	Beaverdam Cr	X	X	X	X	X	X		X
PG-N-117-329-97	Northwest Br	X	X	X	X	X	X		X
PG-N-119-1-94	Oxon Cr	X	X	X	X			X	
PG-N-119-2-94	Oxon Cr	X	X	X	X			X	
PG-N-125-218-97	Beaverdam Cr	X	X	X	X	X	X		X
PG-N-125-228-97	Beaverdam Cr	X	X	X	X	X	X		X
PG-N-130-327-97	Piscataway Cr	X	X	X	X	X	X		X
PG-N-135-231-97	Charles Br	X	X	X	X	X	X		X
PG-N-137-1-94	Indian Cr	X	X	X	X	X		X	

Appendix A (cont.). Summary of the types of data collected at Maryland Biological Stream Survey sites in Prince George's County, 1994-1997. Abbreviations used are as follows: F-IBI - Fish Index of Biotic Integrity; B-IBI - Benthic Macroinvertebrate Index of Biotic Integrity; Fam. IBI - Family-Level Benthic Macroinvertebrate Index of Biotic Integrity; PHI - Physical Habitat Index.

<i>Site</i>	<i>Stream Name</i>	<i>Benthic Macroinvertebrate</i>		<i>Habitat</i>		<i>F-IBI</i>		<i>Fam. IBI</i>	
		<i>Fish</i>	<i>Herpetofauna</i>	<i>Water Chemistry</i>	<i>B-IBI</i>	<i>PHI</i>			
PG-N-259-1-94	Collington Br	X	X	X	X		X	X	
PG-N-259-2-94	Collington Br	X	X	X	X	X	X	X	
PG-N-260-1-94	Indian Cr	X	X	X	X	X			X
PG-N-260-2-94	Indian Cr	X	X	X	X		X	X	
PG-N-271-9-94	Collington Br	X	X	X	X	X	X	X	
PG-N-274-128-97	Honey Br	X	X	X	X	X	X	X	X
PG-S-005-220-95	Mattawoman Creek		X		X		X		
PG-S-007-108-97	Un Trib To Summerville Cr	X	X	X	X		X	X	X
PG-S-032-209-95	Mattawoman Creek	X	X	X	X	X	X	X	X
PG-S-035-301-97	Swanson Cr	X	X	X	X	X	X	X	X
PG-S-045-317-97	Swanson Cr		X		X		X		
PG-S-047-211-97	Rock Br To Spice Cr	X	X	X	X	X	X	X	X
PG-S-052-109-95	Wolf Den Branch	X	X	X	X	X	X	X	X

Appendix B. Physical habitat condition measured by the Maryland Biological Stream Survey, 1994-1997. All variables rated on a scale of 0 (poor) to 20 (optimal) unless otherwise noted.

SUBSTRATE AND INSTREAM COVER

Instream Habitat is rated according to the perceived value of habitat to the fish community. Higher scores are assigned to sites with a variety of habitat types and particle sizes. In addition, higher scores are assigned to sites with a high degree of uneven substrate, including logs and rootwads. In streams where substrate types are favorable but flows are so low that fish are essentially precluded from using the habitat, low scores are assigned. If none of the habitat within a segment is useable by fish, a score of zero is assigned.

Epifaunal Substrate is rated based on the amount and variety of hard, stable substrates usable by benthic macroinvertebrates. Because they inhibit colonization, flocculent materials or fine sediments surrounding otherwise good substrates are assigned low scores. Scores are also reduced when substrates are less stable.

Velocity/Depth Diversity is rated based on the variety of velocity/depth regimes present at a site (slow-shallow, slow-deep, fast-shallow, and fast-deep). As with embeddedness, this metric varies by stream gradient.

Pool/Glide/Eddy Quality is rated based on the variety and spatial complexity of slow or still water habitat within the sample segment. In high-gradient streams, functionally important slow water habitat may exist in the form of larger eddies. Within a category, higher scores are assigned to segments which have undercut banks, woody debris or other types of cover for fish.

Riffle/Run Quality is based on the depth, complexity, and functional importance of riffle/run habitat in the segment, with highest scores assigned to segments dominated by deeper riffle/run areas, stable substrates, and a variety of current velocities.

Embeddedness is a percentage of surface area of larger particles that is surrounded by fine sediments on the stream bottom. In low gradient streams, embeddedness may be high even in relatively unimpaired watersheds.

CHANNEL CHARACTER

Channel Alteration is a measure of large-scale changes in the shape of the stream channel. Channel alteration includes: concrete channels, artificial embankments, obvious straightening of the natural channel, rip-rap, or other structures, as well as recent bar development. Ratings for this metric are based on the presence of artificial structures as well as the existence, extent, and coarseness of point bars, side bars, and mid-channel bars which indicate the degree of flow fluctuations and substrate stability. Evidence of channelization may sometimes be seen in the form of berms that parallel the stream channel.

Bank Stability is rated based on the presence/absence of riparian vegetation and other stabilizing bank materials such as boulders and rootwads, and frequency/size of erosional areas. Sites with steep slopes are not penalized if banks are composed solely of stable materials.

Channel Flow Status is the percentage of the stream channel that has water, with subtractions made for exposed substrates and dewatered areas.

RIPARIAN CORRIDOR

Shading is rated based on estimates of the degree and duration of shading at a site during summer, including any effects of shading caused by land forms.

Appendix B (cont.). Physical habitat condition measured by the Maryland Biological Stream Survey, 1994-1997.
All variables rated on a scale of 0 (poor) to 20 (optimal) unless otherwise noted.

Riparian Buffer is rated according to the size and type of the vegetated riparian buffer zone at the site. Cultivated fields for agriculture that have bare soil to any extent are not considered as riparian buffers. At sites where the buffer width is variable, or direct delivery of storm runoff or sediment to the stream is evident or highly likely, the narrowest representative buffer width in the segment (e.g., 0 if parking lot runoff enters directly to the stream) is measured and recorded even though some of the stream segment may have a well developed riparian buffer.

AESTHETICS/REMOTENESS

Aesthetics are rated according to the visual appeal of the site and presence/absence of human refuse, with highest scores assigned to stream segments with no human refuse and visually outstanding character.

Remoteness is rated based on the absence of detectable human activity and difficulty in accessing the segment.